

ISSN 2224-526X

ҚАЗАҚСТАН РЕСПУБЛИКАСЫ  
ҰЛТТЫҚ ҒЫЛЫМ АКАДЕМИЯСЫНЫҢ

# Х А Б А Р Л А Р Ы

---

---

## ИЗВЕСТИЯ

НАЦИОНАЛЬНОЙ АКАДЕМИИ НАУК  
РЕСПУБЛИКИ КАЗАХСТАН

## NEWS

OF THE NATIONAL ACADEMY OF SCIENCES  
OF THE REPUBLIC OF KAZAKHSTAN

АГРАРЛЫҚ ҒЫЛЫМДАР СЕРИЯСЫ



СЕРИЯ АГРАРНЫХ НАУК



SERIES OF AGRICULTURAL SCIENCES

1 (43)

ҚАҢТАР – АҚПАН 2018 ж.  
ЯНВАРЬ – ФЕВРАЛЬ 2018 г.  
JANUARY – FEBRUARY 2018

2011 ЖЫЛДЫҢ ҚАҢТАР АЙЫНАН ШЫҒА БАСТАҒАН  
ИЗДАЕТСЯ С ЯНВАРЯ 2011 ГОДА  
PUBLISHED SINCE JANUARY 2011

ЖЫЛЫНА 6 РЕТ ШЫҒАДЫ  
ВЫХОДИТ 6 РАЗ В ГОД  
PUBLISHED 6 TIMES A YEAR

АЛМАТЫ, ҚР ҰҒА  
АЛМАТЫ, НАН РК  
ALMATY, NAS RK

Б а с р е д а к т о р

**Есполов Т.И.,**

э.ғ.д, профессор,

ҚР ҰҒА академигі және вице-президенті

Р е д а к ц и я а л қ а с ы:

**Байзақов С.Б.**, э.ғ.д, проф., ҚР ҰҒА академигі (бас редактордың орынбасары); **Тиреуов К.М.**, э.ғ.д, проф., ҚР ҰҒА академигі (бас редактордың орынбасары); **Елешев Р.Е.**, т.ғ.д., проф., ҚР ҰҒА академигі; **Рау А.Г.**, т.ғ.д., проф., ҚР ҰҒА академигі; **Иванов Н.П.**, в.ғ.д, проф., ҚР ҰҒА академигі; **Кешуов С.А.**, т.ғ.д., проф., ҚР ҰҒА академигі; **Мелдебеков А.**, а.ш.ғ.д., проф., ҚР ҰҒА академигі; **Чоманов У.Ч.**, т.ғ.д., проф., ҚР ҰҒА академигі; **Елюбаев С.З.**, а.ш.ғ.д., проф., ҚР ҰҒА академигі; **Садыкулов Т.**, а.ш.ғ.д., проф., академигі; **Баймұқанов Д.А.**, а.ш.ғ.д., проф., ҚР ҰҒА корр-мүшесі; **Сансызбай А.Р.**, а.ш.ғ.д., проф., ҚР ҰҒА корр-мүшесі; **Умбетаев И.**, а.ш.ғ.д., проф., ҚР ҰҒА академигі; **Оспанов С.Р.**, а.ш.ғ.д., проф., ҚР ҰҒА құрметті мүшесі; **Олейченко С.И.**, а.ш.ғ.д., проф.; **Кененбаев С.Б.**, а.ш.ғ.д., проф., ҚР ҰҒА корр-мүшесі; **Омбаев А.М.**, а.ш.ғ.д., проф. ҚР ҰҒА корр-мүшесі; **Молдашев А.Б.**, э.ғ.д., проф., ҚР ҰҒА құрметті мүшесі; **Сагитов А.О.**, б.ғ.д., ҚР ҰҒА академигі; **Сапаров А.С.**, а.ш.ғ.д., проф., ҚР АШҒА академигі; **Балгабаев Н.Н.**, а.ш.ғ.д., проф.; **Умирзаков С.И.**, т.ғ.д, проф.; **Султанов А.А.**, в.ғ.д., проф., ҚР АШҒА академигі; **Алимкулов Ж.С.**, т.ғ.д., проф., ҚР АШҒА академигі; **Сарсембаева Н.Б.**, в.ғ.д., проф.

Р е д а к ц и я к е ñ е с і:

**Fasler-Kan Elizaveta**, Dr., University of asel Switzeland; **Koolmees Petrus Adrianus**, Prof. Dr., Utrecht University, The Netherlands; **Babadoost-Kondri Mohammad**, Prof., University of Illinois, USA; **Yus Aniza Binti Yusof**, Dr., University Putra, Malaysia; **Hesseln Hayley Fawn**, As. Prof., University of Saskatchewan, Canada; **Alex Morgounov**, Pr., International Maize and Wheat Improvement Center Turkey; **Андреш С.**, Молдова Республикасы ҰҒА академигі; **Гаврилюк Н.Н.**, Украина ҰҒА академигі; **Герасимович Л.С.**, Беларусь Республикасының ҰҒА академигі; **Мамедов Г.**, Азербайжан Республикасының ҰҒА академигі; **Шейко И.П.**, Беларусь Республикасының ҰҒА академигі; **Жалнин Э.В.**, т.ғ.д., проф., Ресей; **Боинчан Б.**, а.ш.ғ.д, проф., Молдова Республикасы; **Юлдашбаев Ю.А.**, а.ш.ғ.д, проф., РФА корр-мүшесі, Ресей.

Главный редактор

**Есполов Т.И.,**

доктор эконом. наук, проф.,  
вице-президент и академик НАН РК

Редакционная коллегия:

**Байзаков С.Б.**, доктор эконом. наук, проф., академик НАН РК (заместитель главного редактора); **Тиреуов К.М.**, доктор эконом. наук, проф., академик НАН РК (заместитель главного редактора); **Елешев Р.Е.**, доктор техн. наук, проф., академик НАН РК; **Рау А.Г.**, доктор техн. наук, проф., академик НАН РК; **Иванов Н.П.**, доктор ветеринар. наук, проф., академик НАН РК; **Кешуов С.А.**, доктор техн. наук, проф., академик НАН РК; **Мелдебеков А.**, доктор сельхоз. наук, проф., академик НАН РК; **Чоманов У.Ч.**, доктор техн. наук, проф., академик НАН РК; **Елюбаев С.З.**, доктор сельхоз. наук, проф., академик НАН РК; **Садыкулов Т.**, доктор сельхоз. наук, проф., академик НАН РК; **Баймуқанов Д.А.**, доктор сельхоз. наук, проф., член-корр. НАН РК; **Сансызбай А.Р.**, доктор сельхоз. наук, проф., член-корр. НАН РК; **Умбетаев И.**, доктор сельхоз. наук, проф., академик НАН РК; **Оспанов С.Р.**, доктор сельхоз. наук, проф., Почетный член НАН РК; **Олейченко С.И.**, доктор сельхоз. наук, проф.; **Кененбаев С.Б.**, доктор сельхоз. наук, проф., член-корр. НАН РК; **Омбаев А.М.**, доктор сельхоз. наук, проф. член-корр. НАН РК.; **Молдашев А.Б.**, доктор эконом. наук, проф., Почетный член НАН РК; **Сагитов А.О.**, доктор биол. наук, академик НАН РК; **Сапаров А.С.**, доктор сельхоз. наук, проф., академик АСХН РК; **Балгабаев Н.Н.**, доктор сельхоз. наук, проф.; **Умирзаков С.И.**, доктор техн. наук, проф.; **Султанов А.А.**, доктор ветеринар. наук, проф., академик АСХН РК; **Алимкулов Ж.С.**, доктор техн. наук, проф., академик АСХН РК; **Сарсембаева Н.Б.**, доктор ветеринар. наук, проф.

Редакционный совет:

**Fasler-Kan Elizaveta**, Dr., University of asel Switzeland; **Koolmees Petrus Adrianus**, Prof. Dr., Utrecht University, The Netherlands; **Babadoost-Kondri Mohammad**, Prof., University of Illinois, USA; **Yus Aniza Binti Yusof**, Dr., University Putra, Malaysia; **Hesseln Hayley Fawn**, As.Prof., University of Saskatchewan, Canada; **Alex Morgounov**, Pr., International Maize and Wheat Improvement Center Turkey; **Андреш С.**, академик НАН Республики Молдова; **Гаврилюк Н.Н.**, академик НАН Украины; **Герасимович Л.С.**, академик НАН Республики Беларусь; **Мамедов Г.**, академик НАН Республики Азербайджан; **Шейко И.П.**, академик НАН Республики Беларусь; **Жалнин Э.В.**, доктор техн. наук, проф., Россия; **Боинчан Б.**, доктор сельхоз. наук, проф., Республика Молдова; **Юлдашбаев Ю.А.**, доктор сельхоз. наук, проф., член-корр. РАН, Россия.

**Известия Национальной академии наук Республики Казахстан. Серия аграрных наук.**

**ISSN 2224-526X**

Собственник: ООО «Национальная академия наук Республики Казахстан» (г. Алматы)

Свидетельство о постановке на учет периодического печатного издания в Комитете информации и архивов Министерства культуры и информации Республики Казахстан № 10895-Ж, выданное 30.04.2010 г.

Периодичность 6 раз в год

Тираж: 300 экземпляров

Адрес редакции: 050010, г. Алматы, ул. Шевченко, 28, ком. 219-220, тел. 272-13-19, 272-13-18

<http://nauka-nanrk.kz/agricultural.kz>

---

© Национальная академия наук Республики Казахстан, 2018

Адрес типографии: ИП «Аруна», г. Алматы, ул. Муратбаева, 75

Chief Editor

**Espolov T.I.,**

Dr. economy. Sciences, prof.,  
Vice President and academician of the NAS RK

Editorial Board:

**Baizakov S.B.**, Dr. of economy sciences, prof., academician of NAS RK (deputy editor); **Tireuov K.M.**, Doctor of Economy Sciences., prof., academician of NAS RK (deputy editor); **Eleshev R.E.**, Dr. Of agricultural sciences, prof., academician of NAS RK; **Rau A.G.**, Dr. sciences, prof., academician of NAS RK; **Ivanov N.P.**, Dr. of veterinary sciences, prof., academician of NAS RK; **Keshuov S.A.**, Dr. sciences, prof., academician of NAS RK; **Meldebekov A.**, doctor of agricultural sciences, prof., academician of NAS RK; **Chomanov U.Ch.**, Dr. sciences, prof., academician of NAS RK; **Yelyubayev S.Z.**, Dr. of agricultural sciences, prof., academician of NAS RK; **Sadykulov T.**, Dr. Farm. Sciences, prof., academician of NAS RK; **Baimukanov D.A.**, doctor of agricultural sciences, prof., corresponding member NAS RK; **Sansyzbai A.R.**, doctor of agricultural sciences, prof., corresponding member NAS RK; **Umbetaev I.**, Dr. Farm. Sciences, prof., academician of NAS RK; **Ospanov S.R.**, Dr. agricultural sciences, prof., Honorary Member of NAS RK; **Oleychenko S.N.**, Dr. Of agricultural sciences, prof.; **Kenenbayev S.B.**, Dr. Agricultural sciences, prof., corresponding member NAS RK; **Ombayev A.M.**, Dr. Agricultural sciences, Prof. corresponding member NAS RK; **Moldashev A.B.**, Doctor of Economy sciences, prof., Honorary Member of NAS RK; **Sagitov A.O.**, Dr. biol. sciences, academician of NAS RK; **Saparov A.S.**, Doctor of agricultural sciences, prof., academician of NAS RK; **Balgabaev N.N.**, the doctor agricultural sciences, Prof.; **Umirzakov S.I.**, Dr. Sci. Sciences, Prof.; **Sultanov A.A.**, Dr. of veterinary sciences, prof., academician of the Academy of Agricultural Sciences of Kazakhstan; **Alimkulov J.C.**, Dr. of tekhncial sciences, prof., academician of the Academy of Agricultural sciences of Kazakhstan; **Sarsembayeva N.B.**, Dr. veterinary sciences, prof.

Editorial Board:

**Fasler-Kan Elizaveta**, Dr., University of Basel Switzzeland; **Koolmees Petrus Adrianus**, Prof. Dr., Utrecht University, The Netherlands; **Babadoost-Kondri Mohammad**, Prof., University of Illinois, USA; **Yus Aniza Binti Yusof**, Dr., University Putra, Malaysia; **Hesseln Hayley Fawn**, As. Prof., University of Saskatchewan, Canada; **Alex Morgounov**, candidate of agricultural sciences, International Maize and Wheat Improvement Center Turkey; **Andresh S.**, academician of NAS of Moldova; **Gavriluk N.N.**, academician of NAS of Ukraine; **Gerasimovich L.S.**, academician of NAS of Belorassia; **Mamadov G.**, academician of NAS of Azerbaijan; **Sheiko I.P.**, academician of NAS of Belorassia; **Zhalnin E.V.**, Dr. of technical sciences, professor, Russia, **Boinchan B.**, doctor of agricultural sciences, prof., Moldova; **Yuldashbayev Y.A.**, doctor of agricultural sciences, prof., corresponding member of RAS, Russia.

**News of the National Academy of Sciences of the Republic of Kazakhstan. Series of Agrarian Sciences.**

**ISSN 2224-526X**

Owner: RPA "National Academy of Sciences of the Republic of Kazakhstan" (Almaty)

The certificate of registration of a periodic printed publication in the Committee of Information and Archives of the Ministry of Culture and Information of the Republic of Kazakhstan N 10895-Ж, issued 30.04.2010

Periodicity: 6 times a year

Circulation: 300 copies

Editorial address: 28, Shevchenko str., of.219-220, Almaty, 050010, tel. 272-13-19, 272-13-18,  
<http://nauka-nanrk.kz/> [agricultural.kz](http://agricultural.kz)

---

© National Academy of Sciences of the Republic of Kazakhstan, 2018

Address of printing house: ST "Aruna", 75, Muratbayev str, Almaty

NEWS

OF THE NATIONAL ACADEMY OF SCIENCES OF THE REPUBLIC OF KAZAKHSTAN

SERIES OF AGRICULTURAL SCIENCES

ISSN 2224-526X

Volume 1, Number 43 (2018), 38 – 41

UDC 632.913.1

**B. B. Toyzhigitova<sup>1</sup>, S. Yskak<sup>2</sup>, A. S. Dinasilov<sup>2</sup>, Zh. B. Niyazbekov<sup>2</sup>**

<sup>1</sup>«Kazakh National Agrarian University», Almaty, Kazakhstan,

<sup>2</sup>«Kazakh Research Institute of Protection and Plants Quarantine», Almaty, Kazakhstan.

E-mail: bayan.toyzhigitova.69@mail.ru; sapor1234@mail.ru; alhimzr@mail.ru

**EFFICIENCY OF ENVIRONMENTALLY SAFE TECHNIQUES  
AND METHODS AGAINST THE MELON FLY  
*Myiopardalis pardalina* (Bigot)**

**Abstract.** At the density for mass catching the pheromone glue traps, food baits of 10 pieces/hectare, the significant effect on prevalence of fruits by the melon fly and productivity in the microplot experiments has not been noted. Technologies of pheromone disorientation of males practically do not influence on population of the melon fly due to low efficiency of dispensers. Installation of the pheromone traps in number of 20, 40 and 60 pcs/ha has shown that the quantity of the caught males has averaged 2.05 pcs on 1 trap.

**Keywords:** melon fly, imago, wrecker, pheromone traps, food baits.

**Introduction.** The developed control measures for the melon fly are so far ineffective and suggest spatial isolation of crops, early sowing of early ripening varieties, the use of 3-4-year-old seeds for sowing, control of weeds, removal of afterharvesting residues, fall plowing and a number of other agricultural techniques against the background of an intensive press of insecticide pest. Multiple treatment of crops with insecticides led to the formation of pest resistance to chemical preparations. Alternative methods of control have not yet been developed.

The evaluation of environmentally safe techniques and methods that were used to control other types of fruit flies, their advance to local conditions, as well as their combination with the intelligent use of biologically active substances, pheromones, sterilization and pesticides of the new generation, identification of promising entomophages will allow to develop effective and ecologically friendly methods of controlling the melon fly and to reduce their harmful impact on the environment in integrated melon protection systems in the Republic of Kazakhstan.

In the conditions of Karakalpakstan, bordering of the Kyzylorda region and similar in soil and climatic conditions, based on the study of bioecological features, methods for protecting melons from this pest have been developed with the primary use of agrotechnical and chemical methods of struggle. However, the use of the recommended melon protection scheme does not solve the problem of eliminating the foci of this quarantine pest. Multiple treatment of crops with insecticides with the same active substances (chlorpyrifos, 500 g/l + cypermethrin, 50 g/l) leads to the formation of pest resistance to chemical preparations [1].

The most effective alternatives to the chemical method aimed at suppressing populations of flies, including the melon fly, are analyzed.

**Materials and methods.** Technology of the male vacuum. Mass catch, also called as male vacuum or elimination method, is based on the principle of intensive attraction of males to traps with an attractant. As a result of the removal of males from the population, females remain unfertilized. Mass catch is carried out with the help of a large number of traps, covering as much as possible the zone of action of natural pheromone produced by females of the natural population. On 1 hectare depending on the number of people from 5 to 50 traps are hanged.

When studying the effectiveness of catching flies on pheromone traps, the obtained results were compared with the results of mowing with insect net, 50 double strokes per passage along the diagonal of the field [2, 3].

Technology of pheromone disorientation of males of the melon fly. The method of disorientation: at the beginning of flowering, rubber rings with the content of pheromone (180-200 mg) are hung. Only 500 rings per 1 ha, which corresponds to 90-100 g of preparation per ha. In early summer, pests of the second and third generation (according to observations in control pheromone traps), the same number of rubber rings with pheromone are re-hung. A total of 270-300 g of pheromone is required for protection during the season [4].

Field works were carried out in the South Kazakhstan region and in the entomology laboratory of the plant quarantine department of the KazRIP&PQ.

**Results of the research.** During the growing season, experiments were conducted to determine the effectiveness of individual types of pheromone glue traps, food baits and their optimum density for mass fly catching.

Various types of pheromone traps have been tested: pheromone glue traps with a melon fly dispenser, colored glue traps, food baits in the form of solutions of attracting substances (melon syrup). The traps were set in the period of the mass appearance of the melon fly, from the end of flowering melons, at a distance of 20-30 m from each other in 4-fold repetition. Accounting was conducted every 5 days, taking into account the number of caught individuals and their total number for the season was determined (table 1).

Table 1 – Effectiveness of catching on pheromone, colored and food traps of the melon flies, their effect on melon damage and yield, Maktaaral district, 2017

| Experiment variants        | Frequency | Caught, for the season | Percentage of affected melon fruit | Yielding, c/ha |
|----------------------------|-----------|------------------------|------------------------------------|----------------|
| Pheromone traps, 2 pcs/ha  | 1         | 16                     | 13                                 | 112            |
|                            | 2         | 13                     | 16                                 | 156            |
|                            | 3         | 28                     | 11                                 | 176            |
|                            | 4         | 43                     | 15                                 | 164            |
| Average                    |           | 25.0                   | 13.8                               | 152.0          |
| Pheromone traps, 10 pcs/ha | 1         | 61                     | 12                                 | 167            |
|                            | 2         | 47                     | 9                                  | 157            |
|                            | 3         | 58                     | 3                                  | 183            |
|                            | 4         | 67                     | 11                                 | 158            |
| Average                    |           | 58.3                   | 8.8                                | 166.3          |
| Glue traps, 2 pcs/ha       | 1         | 12                     | 16                                 | 167            |
|                            | 2         | 3                      | 14                                 | 137            |
|                            | 3         | 9                      | 16                                 | 156            |
|                            | 4         | 15                     | 17                                 | 167            |
| Average                    |           | 9.8                    | 15.8                               | 156.8          |
| Glue traps, 10 pcs/ha      | 1         | 16                     | 18                                 | 167            |
|                            | 2         | 17                     | 19                                 | 154            |
|                            | 3         | 15                     | 16                                 | 137            |
|                            | 4         | 14                     | 17                                 | 178            |
| Average                    |           | 15.5                   | 17.5                               | 159.0          |
| Food baits, 2 pcs/ha       | 1         | 18                     | 16                                 | 167            |
|                            | 2         | 7                      | 17                                 | 173            |
|                            | 3         | 6                      | 11                                 | 164            |
|                            | 4         | 7                      | 18                                 | 149            |
| Average                    |           | 9.5                    | 15.5                               | 163.3          |
| Food baits, 10 pcs/ha      | 1         | 32                     | 16                                 | 171            |
|                            | 2         | 14                     | 18                                 | 165            |
|                            | 3         | 26                     | 19                                 | 170            |
|                            | 4         | 17                     | 12                                 | 146            |
|                            |           | 22.3                   | 16.3                               | 163.0          |

The works were conducted in the Maktaaral and Shardara districts of the South Kazakhstan region in stationary fields.

Stationary field N1. Maktaaral district, Kyzylzhar rural district. GPSN 40°51'38.74<sup>11</sup>, E 068°18'24.1<sup>11</sup>, H 250. “Kirov” field.

Stationary field N 2. Maktaaral district, Kyzylzhar rural district. GPSN 40°53'15.1<sup>11</sup>, E 068°18'56,8<sup>11</sup>, H 255. Akimat’s field.

Stationary field N 3. Maktaaral district, Kyzylzhar rural district. GPSN 40°49'29,4<sup>11</sup>, E 068°15'25.8<sup>11</sup>, H 220. Field N3.

Stationary field N4. Shardara district, GPSN 40°53'18,1<sup>11</sup>, E 068°18'56,8<sup>11</sup>, H 309. Field N4.

Stationary field N5. Shardara district, GPSN 40°51'17,2<sup>11</sup>, E 068°21'56,9<sup>11</sup>, H 288. Field N5.

For the primary comparative assessment of the pheromone disorientation of males for the melon fly pest population, the pheromone traps produced in Moldova for trapping the melon fly in the Maktaaral district of the South Kazakhstan region in the amount of 20, 40 and 60 pcs/ha were installed. Accounting conducted on the 30th day after installation showed that the number of trapped males averaged 3.9; 2.7 and 0.9 pcs. per 1 trap, respectively (table 2).

Table 2 – Effectiveness of melon fly catching on pheromone traps (South-Kazakhstan region, Saryagash district, 2017)

| Frequency | Quantity of traps per 1 ha, pcs. |      |      |
|-----------|----------------------------------|------|------|
|           | 20                               | 40   | 60   |
| 1         | 10                               | 11   | 10   |
| 2         | 9                                | 5    | 5    |
| 3         | 9                                | 4    | 5    |
| 4         | 13                               | 9    | 13   |
| Average   | 2.05                             | 0.72 | 0.55 |

Pheromone traps established in the background of a low number of flies and two-fold processing with Angio 247 S.C. in a dosage of 0.15 l/ha showed catch of individual pests. At 20, 40 and 60 traps per 1 hectare, the males of melon fly practically did not fall into traps of white and yellow color with a dispenser.

**Conclusions.** At the density for mass catching with pheromone glue traps, food baits of 10 pcs/ha, there was no significant effect on the affected fruits by melon fly and yield in micro-plot experiments. Technologies of the pheromone disorientation of males have practically no effect on the population of the melon fly due to the low effectiveness of dispensers. The installation of the pheromone traps in the amount of 20, 40 and 60 pcs/ha showed that the number of trapped males averaged 0.55-2.05 pcs on 1 trap.

Thus, the technologies of the pheromone disorientation of males have practically no effect on the population of the melon fly, due to the low effectiveness of the dispensers.

#### REFERENCES

- [1] Torenliyazov E.Sh. et al. Bioecology of development of the melon fly and recommendations for fighting. Ministry of Agriculture and Water Resources of the Republic of Uzbekistan, Nukus Branch of the Tashkent State Agrarian University. – Nukus, 2010. – 32 p. (in Russ.).
- [2] Suleimenov S.I., Abdrakhmanov M.A., Amerguzhin R.Sh., Butaev K.M., Suleimenova Z.Sh., Mukhyshov A.A., Palinka A.P., Tokmurzina Z.Kh., Abdugalieva G.H., Tleubaev N.K. Methodical instructions for the registration and identification of quarantine objects. – Astana, 2009. – 75 p. (in Russ.).
- [3] Fursov V.N. How to collect insect-entomophages // Recommendation. – Kiev: Institute of Zoology, 2003. – 68 p. (in Russ.).
- [4] Lebedeva K.V., Miniyailo V.A., Pyatnova Yu.B. Pheromones of insects. – Moscow: Nauka, 1984. – 269 p. (in Russ.).

Б. Б. Тойжігітова<sup>1</sup>, С. Ыскак<sup>2</sup>, А. С. Динасилов<sup>2</sup>, Ж. Б. Ниязбеков<sup>2</sup>

<sup>1</sup>Қазақ ұлттық аграрлық университеті, Алматы, Қазақстан,

<sup>2</sup>Қазақ өсімдік қорғау және карантин ғылыми-зерттеу институты, Алматы, Қазақстан

### ҚАУЫН ШЫБЫНЫНА ҚАРСЫ ЭКОЛОГИЯЛЫҚ ҚАУПСІЗ ӘДІСТЕРДІ ҚОЛДАНУ ЖӘНЕ ОНЫҢ ТИІМДІЛІГІ *Myiopardalis pardalina* (Bigot)

**Аннотация.** Жаппай аулауда феромон жабысқақ немесе тағамдық тұзақтарға 10 дана/га оңтайлы жиілігі қауын шыбынымен жеміс және өнімнің зақымдану әсері айтарлықтай болған жоқ.

Аталықтарды феромон көмегімен адастыру технологиясы диспенсерлердің тиімділігі төмен болғандықтан іс жүзінде қауын шыбыны популяциясына әсер етпейді.

**Түйін сөздер:** қауын шыбыны, ересек шыбын, зиянкес, феромондық тұтқыш, тағамдық тұтқыш

Б. Б. Тойжігітова<sup>1</sup>, С. Ыскак<sup>2</sup>, А. С. Динасилов<sup>2</sup>, Ж. Б. Ниязбеков<sup>2</sup>

<sup>1</sup>Казахский аграрный университет, Алматы, Казахстан,

<sup>2</sup>ТОО «Казахский научно-исследовательский институт защиты и карантина растений»,  
Алматы, Казахстан

### ЭФФЕКТИВНОСТЬ ЭКОЛОГИЧЕСКИ БЕЗОПАСНЫХ ПРИЕМОМ И МЕТОДОВ ПРОТИВ ДЫННОЙ МУХИ *Myiopardalis pardalina* (Bigot)

**Аннотация.** При плотности для массового отлова феромонных клеевых ловушек, пищевых приманок 10 шт/га существенного влияния на пораженность плодов дынной мухой и урожайность в мелкоделяночных опытах не отмечено. Технологии феромонного дезориентирования самцов практически не оказывают влияния на популяцию дынной мухи вследствие низкой эффективности диспенсеров. Установка феромонных ловушек в количестве 20, 40 и 60 шт./на 1 га показала, что количество отловленных самцов в среднем не превышала 2,05 шт. на 1 ловушку.

**Ключевые слова:** дынная муха, феромонные ловушки, пищевые приманки.

#### About the authors:

Toyzhigitova B.B. – PhD student;

Yskak S. – Candidate of agricul. Sc., head of the expert-scientific sector;

Dinasilov A.S. – Candidate of agricul. Sc., head of the laboratory;

Niyazbekov Zh.B. – Head of Plant Quarantine Department



## **Publication Ethics and Publication Malpractice in the journals of the National Academy of Sciences of the Republic of Kazakhstan**

For information on Ethics in publishing and Ethical guidelines for journal publication see <http://www.elsevier.com/publishingethics> and <http://www.elsevier.com/journal-authors/ethics>.

Submission of an article to the National Academy of Sciences of the Republic of Kazakhstan implies that the described work has not been published previously (except in the form of an abstract or as part of a published lecture or academic thesis or as an electronic preprint, see <http://www.elsevier.com/postingpolicy>), that it is not under consideration for publication elsewhere, that its publication is approved by all authors and tacitly or explicitly by the responsible authorities where the work was carried out, and that, if accepted, it will not be published elsewhere in the same form, in English or in any other language, including electronically without the written consent of the copyright-holder. In particular, translations into English of papers already published in another language are not accepted.

No other forms of scientific misconduct are allowed, such as plagiarism, falsification, fraudulent data, incorrect interpretation of other works, incorrect citations, etc. The National Academy of Sciences of the Republic of Kazakhstan follows the Code of Conduct of the Committee on Publication Ethics (COPE), and follows the COPE Flowcharts for Resolving Cases of Suspected Misconduct ([http://publicationethics.org/files/u2/New\\_Code.pdf](http://publicationethics.org/files/u2/New_Code.pdf)). To verify originality, your article may be checked by the Cross Check originality detection service <http://www.elsevier.com/editors/plagdetect>.

The authors are obliged to participate in peer review process and be ready to provide corrections, clarifications, retractions and apologies when needed. All authors of a paper should have significantly contributed to the research.

The reviewers should provide objective judgments and should point out relevant published works which are not yet cited. Reviewed articles should be treated confidentially. The reviewers will be chosen in such a way that there is no conflict of interests with respect to the research, the authors and/or the research funders.

The editors have complete responsibility and authority to reject or accept a paper, and they will only accept a paper when reasonably certain. They will preserve anonymity of reviewers and promote publication of corrections, clarifications, retractions and apologies when needed. The acceptance of a paper automatically implies the copyright transfer to the National Academy of Sciences of the Republic of Kazakhstan.

The Editorial Board of the National Academy of Sciences of the Republic of Kazakhstan will monitor and safeguard publishing ethics.

Правила оформления статьи для публикации в журнале смотреть на сайте:

[www:nauka-nanrk.kz](http://www.nauka-nanrk.kz)

<http://agricultural.kz/>

Редактор *М. С. Ахметова, Т. М. Апендиев, Д. С. Аленов*  
Верстка на компьютере *Д. Н. Калкабековой*

Подписано в печать 07.02.2018.  
Формат 60x881/8. Бумага офсетная. Печать – ризограф.  
8,25 п.л. Тираж 300. Заказ 1.