ҚАЗАҚСТАН РЕСПУБЛИКАСЫ ҰЛТТЫҚ ҒЫЛЫМ АКАДЕМИЯСЫНЫҢ

ҚАЗАҚ ҰЛТТЫҚ АГРАРЛЫҚ УНИВЕРСИТЕТІ

ХАБАРЛАРЫ

ИЗВЕСТИЯ

НАЦИОНАЛЬНОЙ АКАДЕМИИ НАУК РЕСПУБЛИКИ КАЗАХСТАН

КАЗАХСКИЙ НАЦИОНАЛЬНЫЙ АГРАРНЫЙ УНИВЕРСИТЕТ

NEWS

OF THE NATIONAL ACADEMY OF SCIENCES OF THE REPUBLIC OF KAZAKHSTAN

KAZAKH NATIONAL AGRARIAN UNIVERSITY

АГРАРЛЫҚ ҒЫЛЫМДАР СЕРИЯСЫ ◆ СЕРИЯ АГРАРНЫХ НАУК ◆ SERIES OF AGRICULTURAL SCIENCES

6 (48)

ҚАРАША – ЖЕЛТОҚСАН 2018 ж. НОЯБРЬ – ДЕКАБРЬ 2018 г. NOVEMBER – DECEMBER 2018

2011 ЖЫЛДЫҢ ҚАҢТАР АЙЫНАН ШЫҒА БАСТАҒАН ИЗДАЕТСЯ С ЯНВАРЯ 2011 ГОДА PUBLISHED SINCE JANUARY 2011

> ЖЫЛЫНА 6 РЕТ ШЫҒАДЫ ВЫХОДИТ 6 РАЗ В ГОД PUBLISHED 6 TIMES A YEAR

Басредактор

Есполов Т.И.,

э.ғ.д, профессор, ҚР ҰҒА академигі және вице-президенті

Редакцияалқасы:

Байзаков С.Б., э.ғ.д, проф., ҚР ҰҒА академигі (бас редактордың орынбасары); Тиреуов К.М., э.ғ.д, проф., ҚР ҰҒА академигі (бас редактордың орынбасары); Елешев Р.Е., т.ғ.д., проф., ҚР ҰҒА академигі; Рау А.Г., т.ғ.д., проф., ҚР ҰҒА академигі; Иванов Н.П., в.ғ.д, проф., ҚР ҰҒА академигі; Кешуов С.А., т.ғ.д., проф., ҚР ҰҒА академигі; Мелдебеков А., а.ш.ғ.д., проф., ҚР ҰҒА академигі; Чоманов У.Ч., т.ғ.д., проф., ҚР ҰҒА академигі; Елюбаев С.З., а.ш.ғ.д., проф., ҚР ҰҒА академигі; Садыкулов Т., а.ш.ғ.д., проф., қР ҰҒА корр-мүшесі; Сансызбай А.Р., а.ш.ғ.д., проф., ҚР ҰҒА корр-мүшесі; Умбетаев И., а.ш.ғ.д., проф., ҚР ҰҒА академигі; Оспанов С.Р., а.ш.ғ.д., проф., ҚР ҰҒА күрметті мүшесі; Олейченко С.И., а.ш.ғ.д., проф.; Кененбаев С.Б., а.ш.ғ.д., проф., ҚР ҰҒА корр-мүшесі; Омбаев А.М., а.ш.ғ.д., проф. ҚР ҰҒА корр-мүшесі; Молдашев А.Б., э.ғ.д., проф., ҚР ҰҒА күрметті мүшесі; Сагитов А.О., б.ғ.д., ҚР ҰҒА академигі; Сапаров А.С., а.ш.ғ.д., проф., ҚР АШҒА академигі; Балгабаев Н.Н., а.ш.ғ.д., проф.; Умирзаков С.И., т.ғ.д, проф.; Султанов А.А., в.ғ.д., проф., ҚР АШҒА академигі; Алимкулов Ж.С., т.ғ.д., проф., ҚР АШҒА академигі; Сарсембаева Н.Б., в.ғ.д., проф.

Редакциякенесі:

Fasler-Kan Elizaveta, Dr., University of asel Switzeland; Koolmees Petrus Adrianus, Prof. Dr., Utrecht University, The Netherlands; Babadoost-Kondri Mohammad, Prof., University of Illinois, USA; Yus Aniza Binti Yusof, Dr., University Putra, Malayzia; Hesseln Hayley Fawn, As. Prof., University of Saskatchewan, Canada; Alex Morgounov, Pr., International Maize and Wheat Improvement Center Turkey; Андреш С., Молдова Республикасы ҰҒА академигі; Гаврилюк Н.Н., Украина ҰҒА академигі; Герасимович Л.С., Беларусь Республикасының ҰҒА академигі; Мамедов Г., Азербайджан Республикасының ҰҒА академигі; Шейко И.П., Беларусь Республикасының ҰҒА академигі; Жалнин Э.В., т.ғ.д., проф., Ресей; Боинчан Б., а.ш.ғ.д, проф., Молдова Республикасы; Юлдашбаев Ю.А., а.ш.ғ.д, проф., РҒА корр-мүшесі, Ресей.

Главныйредактор

Есполов Т.И.,

доктор эконом. наук, проф., вице-президент и академик НАН РК

Редакционнаяколлегия:

Байзаков С.Б., доктор эконом. наук, проф., академик НАН РК (заместитель главного редактора); Тиреуов К.М., доктор эконом. наук., проф., академик НАН РК (заместитель главного редактора); Елешев Р.Е., доктор техн. наук, проф., академик НАН РК; Рау А.Г., доктор техн. наук, проф., академик НАН РК; Иванов Н.П., доктор ветеринар. наук, проф., академик НАН РК; Кешуов С.А., доктор техн. наук, проф., академик НАН РК; Мелдебеков А., доктор сельхоз. наук, проф., академик НАН РК; Садыкулов Т., доктор сельхоз. наук, проф., академик НАН РК; Садыкулов Т., доктор сельхоз. наук, проф., академик НАН РК; Баймуканов Д.А., доктор сельхоз. наук, проф., член-корр. НАН РК; Сансызбай А.Р., доктор сельхоз. наук, проф., член-корр. НАН РК; Умбетаев И., доктор сельхоз. наук, проф., академик НАН РК; Олейченко С.И., доктор сельхоз. наук, проф.; Кененбаев С.Б., доктор сельхоз. наук, проф., член-корр. НАН РК; Омбаев А.М., доктор сельхоз. наук, проф.; Кененбаев С.Б., доктор сельхоз. наук, проф., член-корр. НАН РК; Омбаев А.М., доктор сельхоз. наук, проф., доктор эконом. наук, проф., Почетный член НАН РК; Сагитов А.О., доктор биол. наук, академик НАН РК; Сапаров А.С., доктор сельхоз. наук, проф., академик АСХН РК; Балгабаев Н.Н., доктор сельхоз. наук, проф.; Умирзаков С.И., доктор техн. наук, проф., академик АСХН РК; Сарсембаева Н.Б., доктор ветеринар. наук, проф.

Редакционный совет:

Fasler-Kan Elizaveta, Dr., University of asel Switzeland; Koolmees Petrus Adrianus, Prof. Dr., Utrecht University, The Netherlands; Babadoost-Kondri Mohammad, Prof., University of Illinois, USA; Yus Aniza Binti Yusof, Dr., University Putra, Malayzia; Hesseln Hayley Fawn, As.Prof., University of Saskatchewan, Canada; Alex Morgounov, Pr., International Maize and Wheat Improvement Center Turkey; Андреш С., академик НАН Республики Молдова; Гаврилюк Н.Н., академик НАН Украины; Герасимович Л.С., академик НАН Республики Беларусь; Мамедов Г., академик НАН Республики Азербайджан; Шейко И.П., академик НАН Республики Беларусь; Жалнин Э.В., доктор техн. наук, проф., Россия; Боинчан Б., доктор сельхоз. наук, проф., Республика Молдова; Юлдашбаев Ю.А., доктор сельхоз. наук, проф., член-корр. РАН, Россия.

Известия Национальной академии наук Республики Казахстан. Серия аграрных наук. ISSN 2224-526X

Собственник: РОО «Национальная академия наук Республики Казахстан» (г. Алматы) Свидетельство о постановке на учет периодического печатного издания в Комитете информации и архивов Министерства культуры и информации Республики Казахстан № 10895-Ж, выданное 30.04.2010 г.

Периодичность 6 раз в год Тираж: 300 экземпляров

Адрес редакции: 050010, г. Алматы, ул. Шевченко, 28, ком. 219-220, тел. 272-13-19, 272-13-18

http://nauka-nanrk.kz/agricultural.kz

© Национальная академия наук Республики Казахстан, 2018

Адрес типографии: ИП «Аруна», г. Алматы, ул. Муратбаева, 75

ChiefEditor

Espolov T.I.,

Dr. economy. Sciences, prof., Vice President and academician of the NAS RK

Editorial Board:

Baizakov S.B., Dr. of economy sciences, prof., academician of NAS RK (deputy editor); Tireuov K.M., Doctor of Economy Sciences., prof., academician of NAS RK (deputy editor); Eleshev R.E., Dr. Of agricultural sciences, prof., academician of NAS RK; Rau A.G., Dr. sciences, prof., academician of NAS RK; Ivanov N.P., Dr. of veterinary sciences, prof., academician of NAS RK; Keshuov S.A., Dr. sciences, prof., academician of NAS RK; Meldebekov A., doctor of agricultural sciences, prof., academician of NAS RK; Chomanov U.Ch., Dr. sciences, prof., academician of NAS RK; Yelvubayev S.Z., Dr. of agricultural sciences, prof., academician of NAS RK; Sadykulov T., Dr. Farm. Sciences, prof., academician of NAS RK; Baimukanov D.A., doctor of agricultural sciences, prof., corresponding member NAS RK; Sansyzbai A.R., doctor of agricultural sciences, prof., corresponding member NAS RK; Umbetaev I., Dr. Farm. Sciences, prof., academician of NAS RK; Ospanov S.R., Dr. agricultural sciences, prof., Honorary Member of NAS RK; Oleychenko S.N., Dr. Of agricultural sciences, prof.; Kenenbayev S.B., Dr. Agricultural sciences, prof., corresponding member NAS RK; Ombayev A.M., Dr. Agricultural sciences, Prof. corresponding member NAS RK; Moldashev A.B., Doctor of Economy sciences, prof., Honorary Member of NAS RK; Sagitov A.O., Dr. biol. sciences, academician of NAS RK; Saparov A.S., Doctor of agricultural sciences, prof., academician of NAS RK; Balgabaev N.N., the doctor agricultural sciences, Prof.; Umirzakov S.I., Dr. Sci. Sciences, Prof.; Sultanov A.A., Dr. of veterinary sciences, prof., academician of the Academy of Agricultural Sciences of Kazakhstan; Alimkulov J.C., Dr. of tekhnical sciences, prof., academician of the Academy of Agricultural sciences of Kazakhstan; Sarsembayeva N.B., Dr. veterinary sciences, prof.

Editorial Board:

Fasler-Kan Elizaveta, Dr., University of Basel Switzeland; Koolmees Petrus Adrianus, Prof. Dr., Utrecht University, The Netherlands; Babadoost-Kondri Mohammad, Prof., University of Illinois, USA; Yus Aniza Binti Yusof, Dr., University Putra, Malayzia; Hesseln Hayley Fawn, As. Prof., University of Saskatchewan, Canada; Alex Morgounov, candidate of agricultural sciences, International Maize and Wheat Improvement Center Turkey; Andresh S., academician of NAS of Moldova; Gavriluk N.N., academician of NAS of Ucraine; Gerasimovich L.S., academician of NAS of Belorassia; Mamadov G., academician of NAS of Azerbaijan; Sheiko I.P., academician of NAS of Belorassia; Zhalnin E.V., Dr. of technical sciences, professor, Russia, Boinchan B., doctor of agricultural sciences, prof., Moldova; Yuldashbayev Y.A., doctor of agricultural sciences, prof., corresponding member of RAS, Russia.

News of the National Academy of Sciences of the Republic of Kazakhstan. Series of Agrarian Sciences. ISSN 2224-526X

Owner: RPA "National Academy of Sciences of the Republic of Kazakhstan" (Almaty)

The certificate of registration of a periodic printed publication in the Committee of Information and Archives of the Ministry of Culture and Information of the Republic of Kazakhstan N 10895-W, issued 30.04.2010

Periodicity: 6 times a year Circulation: 300 copies

Editorial address: 28, Shevchenko str., of.219-220, Almaty, 050010, tel. 272-13-19, 272-13-18,

http://nauka-nanrk.kz / agricultural.kz

© National Academy of Sciences of the Republic of Kazakhstan, 2018

Address of printing house: ST "Aruna", 75, Muratbayev str, Almaty

____ 4 ____

NEWS

OF THE NATIONAL ACADEMY OF SCIENCES OF THE REPUBLIC OF KAZAKHSTAN SERIES OF AGRICULTURAL SCIENCES

ISSN 2224-526X

Volume 6, Number 48 (2018), 15 – 19

https://doi.org/10.32014/2018. 2224-526X.13

UDC 663.433

S. M. Shintassova¹, G. I. Baigaziyeva¹, T. F. Kiseleva²

¹Almaty Technological University, Almaty, Kazakhstan, ²Kemerovo State University, Kemerovo, Russia. E-mail: saida_atu@mail.ru, bgulgaishailias@mail.ru, kisseleva.tf@mail.ru,

INTERRELATION OF THE CRITERIA OF BEER WORT COLOR GRADE AND BARLEY GRAIN CONTAMINATION RATE

Abstract. There is an interconnection between grain color grade and contamination rate. The purpose of this work is to research the influence of ion-ozonic explosive cavitation on barley grain contamination and wort color grade. It has been revealed that brewing barley grains without any treatment have the first degree of contamination, and the wort received from epy malt of these grains does not correspond to the quality regarding indicators of transparency and color grade. Meanwhile, the treated samples had no degree of contamination and the wort conformed to the requirements of the standard.

Keywords: grain, barley, wort, contamination, transparency, color grade, ion-ozonic explosive cavitation.

Introduction. The main thing in barley grain production is high-quality treatment, receiving an environmentally friendly product, high-quality storage, barley seed processing into malt and beer with prevention of distribution and activity of the corresponding microorganisms in them [1]. Mites are the most widespread pest of brewing barley which sizes do not exceed 1 mm. Mites like heat and moisture, favorable temperature for their reproduction is 18-30 °C. For brewing, only the first class of contamination is admissible, and they try to use such grain as soon as possible.

Also, the weevil that eats the inside of the grain is very dangerous to barley, and any degree of contamination evidences about unfitness of barley for malting [2]. Body length of weevils together with the proboscis is from 3 to 5 mm. The linear sizes of grain microflora (viruses, yeast, bacteria, mold fungi) are from 100 nanometers (nm) and above. The sizes of insect pests make from 1 000 000 to 10 000 000 nm [3].

Historically, bactericidal treatment at processing enterprises generally is carried out by chemical reagents [4]. For example, an antioxidant – ascorbic acid, which slows down the growth of grain pathogens, is used as a chemical reagent [5].

Using such substances often makes an undesirable impact on the product; its taste, pH and other parameters change; it can be followed by emergence of cancerogenic compounds. For a long time there was no alternative to use of chemicals [4].

Now the modern science and practice has a wide range of ways of treatment. One of such ways is the use of radiation. Different sources of radiation are used for radiation of foodstuff: radioisotopes (cobalt-60 or caesium-137) and accelerators of electrons. Radiation leads to suppression of activity or death of insects and pathogenic microorganisms, and that gives the chance to increase safety of products and to increase the storage periods of substances [6-8].

Modern methods of grain disinfecting during malting also include the promising method of processing in an ultrahigh frequency electromagnetic field [9].

The research in the field of microbiology should be also noted: scientists from Germany have suggested using lactic bacteria as a starting culture for improvement of biological and technological properties of barley malt. All treated barley samples have shown considerable decrease in aerobic bacteria

(by 99,8 %) and stimulation of growth of yeast in comparison with untreated control [10]. In Thailand, Bacillus megaterium powder, which is dried up by dispersion, is intended for fight against the diseases affecting rice. Powdered skim milk which is a part of this powder keeps viability of cells, protecting the cells from a sharp loss of free water and a change of temperature during storage [11].

Formation of beer mash color grade is closely connected with the degree of contamination of barley grain. Color grade is one of the most important parameters of brewing malt quality [12]. Color grade of wort indicates the quality of malt from the perspective of its suitability for production of light beer. In the «color grade» concept, not only the characteristic shade is important, but also transparency, existence or lack of color range [2].

Ion-ozonic explosive cavitation is a new step to effective, environmentally friendly, reliable production free from chemical disinfectants.

Death of bacteria happens due to oxidation of proteinaceous and lipidic structures, and also other inclusions of colloidal appearances which form the external cover and the cytoplasmatic membrane. Charged particles, getting in a biological cell of a bacterium, break the cell's energy balance, i.e. doom the bacterium to unproductive activity, and the bacterium perishes.

Death of viruses during ion-ozonic explosive cavitation happens due to oxidation of the protective protein coat of nucleic acids which form living cells of viruses [13].

Objects and methods of research. We have investigated the brewing barley of Sanshain grade, German selection, but which had been grown up in Kazakhstan and used for production of malt. Sampling and laboratory analyses were carried out according to the requirements of State standard:

- 1) barley for brewing in accordance with State standard 5060-86 «Barley for brewing. Specifications» [14];
- 2) light barley malt in accordance with State standard 29294-92 «Brewing barley malt. Specifications» [15].

Contamination by pests in accordance with State standard 13586-4-83 «Grain. Methods for determination of infested grain and its damage» [16], using the corresponding screen set (the top screen with openings of 2,5 mm, the lower one – with openings of 1,5 mm). 1 kg of barley was screened using the screens during 2 minutes. If grain temperature was lower than 5 °C, the received screenings and the undersize were warmed up to the temperature of 25-30 °C, until insects begun to move, then counted the number of weevils and mites [17].

Also, the degree of contamination was controlled by means of the Indicator of grain and grain products contamination by pests of IZS type representing «Lozar» set of special traps and bars for their immersion. In the store of traps, they placed a food bait for insects who, in turn, attracted by the smell of baits, got through openings to the store and remained in it. Then, after a certain period of time, they took the traps out from the bulk-grain and defined the number and the species composition of insects [18].

Color grade and transparency of wort were determined with the use of a spectrophotometer. Measurement of optical density was made at 430 nm, and color grade in ABS units was received by multiplication of optical density by the known coefficient. The wort was clarified before the analysis. If wort turbidity was higher than 1 ABS unit, it was filtered to full transparency. To define whether the studied sample is transparent enough, they measured its optical density at 700 and 430 nm (A_{700} and A_{430}). If $A_{700} \le 0.039 A_{430}$, then the sample was considered transparent. Color grade was defined by measuring optical density at 430±0,5 nm in relation to water in ditches 5 or 10 mm thick.

Wort was diluted so that optical density would be within the linear dependence field «optical density – malt concentration» (value to 0,8). Preliminarily, they checked optical density on water in the spectrophotometer which had to make 0,00.

Color grade (C, in ABC units) was calculated using the following formula:

$$C = A_{430} F \cdot 25$$
,

where A_{430} – light absorption at 430 nm in a 10 mm thick ditch; 25 – coefficient; F – dilution factor [19].

Barley was treated using an ion-ozonic explosive cavitation installation. Input parameters were as follows: ozone concentration 2, 4, 6 mg/m³, ion concentration – 500±20, 50250±250, 100000±25 un/cm³ (ion-ozonic mix), explosive cavitation – 2, 4, 6 at, exposure of 5, 10, 15 minutes.

The accepted plan of active planning of the experiment has allowed to receive the influence of three factors on grain disinfecting under eight options of experience. Data processing was made by means of Mathcad 140 program.

Results and their discussion. Grain contamination is important as it can result in possible risks for health in cases of consumption of polluted grain in human foodstuff [20]. The purpose of comprehensive protection of plants is to consider all available methods which suppress the development of harmful organisms. Many methods of indirect protection can be more effective and cheap than use of pesticides [21]. Ion-ozonic explosive cavitation treatment has shown its efficiency during our research.


When studying barley contamination by insect pests, it has been revealed that the number of mites and weevils after grain treatment by ion-ozonic explosive cavitation equals to zero, while the control sample had the first degree of contamination (2 weevils and 15 mites) [22].

It should be noted that scientists from China investigated the influence of high hydrostatic pressure treatment on physical-chemical and sensory characteristics of wheat beer during which its impact on microorganisms has been studied. It has been revealed that an alternative to heat treatment is high hydrostatic pressure treatment which can effectively lead to destruction of microorganisms in order to increase safety and to extend storage periods of foodstuff [23].

The color grade indicator of laboratory wort can also characterize the quality of soluble malt. During investigation of wort color grade (*l*) with the use of Mathcad 140 program, it has been revealed that the entered parameters allowed receiving a mathematical functional dependence of this indicator with the analytical expression describing this criterion [24]:

$$ffl(w):=d_0+d_1w^2+d_2w^3+d_3w^3+d_4w^4+d_5w^5+d_6w^6+d_7w^7$$

As a result, we received a schedule (figure) constructed on experimental points: l = l.



The diagram, constructed on experimental points by determination of color grade criterion

Measurement of color of the wort received from malt is one of the most important quality parameters which are subject to control [25]. Superior quality light malt has to have color grade no more than 0,18 ABS. According to our research, color grade of the laboratory wort control sample makes 1,687 ABS; that of test samples processed at high parameters – 0,200-0,240, and that of samples with low parameters of processing – 0,028-0,075 ABS. As this research has shown, ion-ozonic explosive cavitation exerts positive impact not only on the color grade of laboratory wort, but also on its transparency. The wort prepared from processed malt samples was transparent, unlike the control sample which had slight opalescence.

It can be connected with the fact that, the control sample had the first degree of infection during determination of contamination by pests. It is known that wort transparency depends on microbiological quality of barley. Pollution of barley by fungi, insects, and bacteria results in turbidity of the initial wort [25].

Conclusion. The received results show that, within the range of the set processing parameters (ozone concentration of 2, 4, 6 mg/m³, that of ion -500 ± 20 , 50250 ± 250 , 100000 ± 25 un/cm³, explosive cavitation -2, 4, 6 at, time -5, 10, 15 minutes), insect pests, their larvae, various viruses and bacteria perish.

It is necessary to emphasize that molecular ions of air oxygen are activated, and atomic ions are suppressed, i.e. a weak or sick cell is affected. Therefore there is an increase in epy biological value of grain products. Ozone is a powerful oxidizer – it is the strongest disinfectant in nature. Ozonization destroys everything, including pathogenic bacteria, viruses and microorganisms. Ozone dissolved in water has a preserving effect. Thus, comprehensive ion-ozonic processing of grain in the cavitation zone has a disinfecting effect, destroys insect pests, increases the product's biological value, increases stability against external influences, promotes improvement of viability, and reduces humidity by 2-3 %. Cavitation itself strengthens the positive influence of ion-ozonic processing on grain properties [26].

Theoretical justification of experimental studies shows also that, as a result of destruction (oxidation) of coloring agents, the criterion of color grade has changed to the norm – from 1,687 to 0,028 %.

The research results have shown that, as a result of treatment, the best concentration of ozone when determining the criterion of color grade was 2 mg/m^3 , that of molecular ions -500 un/cm^3 , excessive pressure during cavitation corresponded to 2 at, the best exposure time -5 minutes.

Ion-ozonic explosive cavitation, having an efficient sterilizing effect, allows reducing expenses on disinfecting substances, reduces consumption of energy resources, promotes obtaining social benefit and profit arising due to creation of a qualitative, safe and environmentally friendly ready-made product of increased biological value [27].

REFERENCES

- [1] Maemerov M.M., Iztayev A.I., Kulazhanov T.K., Iskakova G.K. (2011) Scientific bases of ion-ozonic technology of grain and grain products treatment, monograph. 246 p. Aleiron, Almaty. ISBN: 978-601-239-187-9.
- [2] Khosiyev O.A., Khosiyev A.M., Tsugkiyeva V.B. (2016) Brewing Technology. 560 p. Lan, St. Petersburg, Moscow, Krasnodar. ISBN: 978-5-8114-1224-2.
- [3] Dzhankurazov B.O., Dzhankurazov K.B. (2013) To protect the golden grain of Kazakhstan. 165 p. Aleiron, Almaty. ISBN: 9965-476-53-5.
 - [4] Krasnochub A.V. (2003) Ensuring microbiological cleanliness at food productions, Beer and drinks, 5, 30-31 p.
- [5] Mina U., Fuloria A., Aggarwal R. (2016) Effect of ozone and antioxidants on wheat and its pathogen Bipolaris soro-kininana, Cer. Res. Com. 44 (4), 594-604 p. https://dx.doi.org/10.1556/0806.44.2016.039.
- [6] Mussina O.N., Konovalov K.L. (2016) Radiation treatment of food staples and foodstuff by ionizing radiation, Food industry, 8, 46-49 p.
 - [7] Nosova T., Burkhanova M. (2017), Radiated products, Newspaper «I am a buyer and owner». 7, 13.
- [8] Rozhdestvenskaya L.N., Bryazgin A.A., Korobeynikov M.V. (2016) Prerequisites and justifications of ionizing radiation for treatment of food products, Food industry, 11, 39-45 p.
- [9] KretovaYu.I., Cherkassova E.I., Cherkassova M.O. (2014) The method of microwave field energy in ensuring quality of vegetable food staples and foodstuff, the Collection of scientific articles following the results of the All-russian scientific and practical conference with the international participation «Science and practice in quality management, metrology and certification», Moscow, 123-126 p.
- [10] Lorenzo C. Peyer, Marcel De Kruijf, Jim O'Mahony, Lorenzo De Colli, Martin Danaher, Martin Zarnkow, Fritz Jacob, Elke K. Arendt. (2017) Lactobacillus brevis R2Δ as starter culture to improve biological and technological qualities of barley malt, Eur. Food Res. Technol. 243 (8), 1363-1374 p. https://dx.doi.org/10.1007/s00217-017-2847-9.
- [11] Chumthong A., Wiwattanapatapee R., Viernstein H., Pengnoo A., Kanjanamaneesathian M. (2016) Spray-dried powder of Bacillus megaterium for control of rice sheath blight disease: Formulation protocol and efficacy testing in laboratory and greenhouse, Cer. Res. Com. 44 (6), 131-140 p. https://dx.doi.org/10.1556/0806.43.2015.034.
 - [12] Yermolayev S.V. (2015) Formation of color grade in caramel malt, Beer and drinks, 2, 24-26 p.
- [13] Maemerov M.M., Kulazhanov K.S., Iztayev A.I. (2001) Ion-ozon technology in the production of grain products. 213 p. NITS «Gylym», Almaty. ISBN: 9965-07-053-9.
- [14] State standard 5060-86 «Barley for brewing. Specifications». Introduction 1988-07-01. Moscow: Publishing house of standards, 1986. 5 p.
- [15] State standard 29294-92 Brewing barley malt. Specifications. Introduction 1993-06-01. Moscow: Publishing house of standards, 1992. 19 p.
- [16] State standard 13586-4-83 Grain. Methods for determination of infested grain and its damage. Introduction 1984-07-01. Moscow: Publishing house of standards, 1983. 4 p.

- [17] Muravitskaya L.V. (1987) Technical and chemical control of brewing and nonalcoholic productions and basis of product quality control. 256 p. Agropromizdat, Moscow.
- [18] Feidengold V.B., Temirbekova S.A. (2014) Laboratory equipment for quality control of grain and products, resulted from its processing. 248 p. DeLiplus, Moscow. ISBN: 978-5-905170-57-7.
- [19] Danina M.M. (2013) Research methods of properties of raw materials, fermentation products and alcohol-free drinks. Laboratory works. P. 13-14, National Research University of Information Technologies, Mechanics and Optics (NRU ITMO), Saint Petersburg.
- [20] Popovski, S., Kos, K., Strajn, B. Jakovac, Celar, F. A. (2017) Fusarium spp. incidence and DON Contamination in different wheat varieties correlated with the environmental factors, Cer. Res. Com. 45 (1), 114-123 p. https://dx.doi.org/10.1556/0806.44.2016.043.
- [21] Matusinsky, P., Vanova, M., Tvaruzek, L., Polisenska, I., Janecek, M., Smutny, V. (2016) Soil management technologies and mycotoxin contamination of wheat and barley grain, Cer. Res. Com. 44 (2), 320-329 p. https://dx.doi.org/10.1556/0806.44.2016.021.
- [22] Shintassova S., Baigaziyeva G. (2017) Application of innovative technologies in the production of malting, Proceedings of International conference on engineering research and applications (ICERA-17), Istanbul, Turkey, 319-322 p.
- [23] Hua Yin, Jianjun Dong, Junhong Yu, Zongming Chang, Zhonghua Qian, Mingli Liu, Shuli Huang, Xiaocong Hu, Xiaolin Liu, Yang Deng and Dongfeng Wang. (2016) A preliminary study about the influence of high hydrostatic pressure processing on the physicochemical and sensorial properties of a cloudy wheat beer, J. Inst. Brew. 122 (3), 462-467 p. https://dx.doi.org/10.1002/jib.344.
- [24] Lifanchikov A.N., Voskoboynik M. P. (1975) Economic-mathematical methods of planning of food industry development. 248 p. Pishchevaya promyshlennost, Moscow.
- [25] Giuseppe Perretti, Simona Floridi, Benedetta Turchetti, Ombretta Marconi and Paolo Fantozzi. (2012) Quality Control of Malt: Turbidity Problems of Standard Worts Given by the Presence of Microbial Cells, J. Inst. Brew. 117 (2), 212-216 p. https://dx.doi.org/10.1002/j.2050-0416.2011.tb00463.x.
- [26] Iztayev A.I., Kulazhanov T.K., Maemerov M.M., Asangalieyva Zh.R., Iztayev B.A., Sarlybayeva L.M. (2015) Electrical and physical methods of treatment of a grain at elevators and cereal-processing enterprises. 172 p. LLP «Publishing house LEM», Almaty.
- [27] Shintassova S.M., Maemerov M.M., Baigaziyeva G.I., Moldakarimov A.A. (2017) Ion-ozonic explosive cavitation is an ecologically and biologically pure factor of malting intensification, Proceedings of international scientific and practical conference «Innovative development of food, light and hospitality industry», Almaty: ATU. P. 23-24.

С. М. Шинтасова¹, Г. И. Байгазиева¹, Т. Ф. Киселева²

¹Алматы технологиялық университеті, Алматы, Қазақстан, ²Кемерово мемлекеттік университеті, Кемерово, Ресей

АРПА ДӘНДЕРІНІҢ ЗАРЯДТАЛУ ДӘРЕЖЕСІ МЕН СЫРА АШЫТҚЫСЫНЫҢ ТҮСТІЛІКТЕРІ КРИТЕРИЙЛЕРІНІҢ ӨЗАРА БАЙЛАНЫСЫ

Аннотация. Дәндердің түстілігі мен зарядтылығы арасында байланыс болады. Осы жұмыстың мақсаты болып арпа дәні мен сыра ашытқысының зарядтылығына иондыозондық жарылыстыкавитациялық әсерін зерттеу табылады. Сыра қайнататын арпа дәнінің қандай да бір өңдеусіз зарядтылықтың бірінші дәрежесінде болатындығы, ал осы дәннің уытынан алынған сыра ашытқысы сапа жағынан түстілік пен мөлдірлік көрсеткіштеріне сәйкес келмейтіндігі анықталды. Ал өңделген үлгілерде зарядтылықтың дәрежесі болмаған және сыра ашытқысы стандарт талаптарына сәйкес келеді.

Түйін сөздер: астық, арпа, сыра ашытқысы, зарядтылық, мөлдірлік, түстілік, иондыозондық жарылыстыкавитациялық.

С. М. Шинтасова¹, Г. И. Байгазиева¹, Т. Ф. Киселева²

¹ Алматинский технологический университет, Алматы, Казахстан, ² Кемеровский государственный университет, Кемерово, Россия

ВЗАИМОСВЯЗЬ КРИТЕРИЕВ ЦВЕТНОСТИ ПИВНОГО СУСЛА СО СТЕПЕНЬЮ ЗАРАЖЕННОСТИ ЗЕРНА ЯЧМЕНЯ

Аннотация. Существует связь между цветностью и зараженностью зерна. Целью данной работы являлось исследование влияния ионоозонной взрывокавитации на зараженность зерна ячменя и цветность сусла. Выявлено, что зерно пивоваренного ячменя без какой-либо обработки имеет первую степень заражения, а сусло, полученное из солода данного зерна, не соответствует качеству по показателям прозрачности и цветности. В то время как обработанные образцы не имели степени заражения и сусло соответствовало требованиям стандарта.

Ключевые слова: зерно, ячмень, сусло, зараженность, прозрачность, цветность, ионоозонная взрывокавитация.

Information about authors:

Shintassova Saida Muradovna, Almaty Technological University, department «Technology of bread products and processing plants», Master of Engineering, PhD candidate; saida_atu@mail.ru; https://orcid.org/0000-0001-6269-4675

Baigaziyeva Gulgaisha Ilyasovna, Almaty Technological University, department «Technology of bread products and processing plants», Ph.D. in Biology, associate professor; bgulgaishailias@mail.ru; https://orcid.org/0000-0003-4860-9918

Kiseleva Tatyana Fyodorovna, Kemerovo State University, department «Technology of Fermentation and Canning», D. Sc. in Engineering, professor; kisseleva.tf@mail.ru; https://orcid.org/0000-0003-1886-3544

Publication Ethics and Publication Malpractice in the journals of the National Academy of Sciences of the Republic of Kazakhstan

For information on Ethics in publishing and Ethical guidelines for journal publication see http://www.elsevier.com/publishingethics and http://www.elsevier.com/journal-authors/ethics.

Submission of an article to the National Academy of Sciences of the Republic of Kazakhstan implies that the described work has not been published previously (except in the form of an abstract or as part of a published lecture or academic thesis or as an electronic preprint, see http://www.elsevier.com/postingpolicy), that it is not under consideration for publication elsewhere, that its publication is approved by all authors and tacitly or explicitly by the responsible authorities where the work was carried out, and that, if accepted, it will not be published elsewhere in the same form, in English or in any other language, including electronically without the written consent of the copyright-holder. In particular, translations into English of papers already published in another language are not accepted.

No other forms of scientific misconduct are allowed, such as plagiarism, falsification, fraudulent data, incorrect interpretation of other works, incorrect citations, etc. The National Academy of Sciences of the Republic of Kazakhstan follows the Code of Conduct of the Committee on Publication Ethics (COPE), and follows the COPE Flowcharts for Resolving Cases of Suspected Misconduct (http://publicationethics.org/files/u2/New_Code.pdf). To verify originality, your article may be checked by the Cross Check originality detection service http://www.elsevier.com/editors/plagdetect.

The authors are obliged to participate in peer review process and be ready to provide corrections, clarifications, retractions and apologies when needed. All authors of a paper should have significantly contributed to the research.

The reviewers should provide objective judgments and should point out relevant published works which are not yet cited. Reviewed articles should be treated confidentially. The reviewers will be chosen in such a way that there is no conflict of interests with respect to the research, the authors and/or the research funders.

The editors have complete responsibility and authority to reject or accept a paper, and they will only accept a paper when reasonably certain. They will preserve anonymity of reviewers and promote publication of corrections, clarifications, retractions and apologies when needed. The acceptance of a paper automatically implies the copyright transfer to the National Academy of Sciences of the Republic of Kazakhstan.

The Editorial Board of the National Academy of Sciences of the Republic of Kazakhstan will monitor and safeguard publishing ethics.

Правила оформления статьи для публикации в журнале смотреть на сайте:

www:nauka-nanrk.kz http://agricultural.kz/

Редактор М. С. Ахметова, Т. М. Апендиев, Д. С. Аленов Верстка на компьютере Д. Н. Калкабековой

Подписано в печать 30.11.2018. Формат 60х881/8. Бумага офсетная. Печать – ризограф. 6,25 п.л. Тираж 300. Заказ 6.