

ISSN 2224-526X

ҚАЗАҚСТАН РЕСПУБЛИКАСЫ
ҰЛТТЫҚ ҒЫЛЫМ АКАДЕМИЯСЫНЫҢ

ҚАЗАҚ ҰЛТТЫҚ АГРАРЛЫҚ УНИВЕРСИТЕТИ

Х А Б А Р Л А Р Ы

ИЗВЕСТИЯ

НАЦИОНАЛЬНОЙ АКАДЕМИИ НАУК
РЕСПУБЛИКИ КАЗАХСТАН

КАЗАХСКИЙ НАЦИОНАЛЬНЫЙ
АГРАРНЫЙ УНИВЕРСИТЕТ

NEWS

OF THE NATIONAL ACADEMY OF SCIENCES
OF THE REPUBLIC OF KAZAKHSTAN

KAZAKH NATIONAL
AGRARIAN UNIVERSITY

АГРАРЛЫҚ ҒЫЛЫМДАР СЕРИЯСЫ



СЕРИЯ АГРАРНЫХ НАУК



SERIES OF AGRICULTURAL SCIENCES

5 (47)

ҚЫРКҮЙЕК – ҚАЗАН 2018 ж.
СЕНТЯБРЬ – ОКТЯБРЬ 2018 г.
SEPTEMBER – OCTOBER 2018

2011 ЖЫЛДЫҢ ҚАҢТАР АЙЫНАН ШЫҒА БАСТАҒАН
ИЗДАЕТСЯ С ЯНВАРЯ 2011 ГОДА
PUBLISHED SINCE JANUARY 2011

ЖЫЛЫНА 6 РЕТ ШЫҒАДЫ
ВЫХОДИТ 6 РАЗ В ГОД
PUBLISHED 6 TIMES A YEAR

Б а с р е д а к т о р

Есполов Т.И.,

э.ғ.д, профессор,

ҚР ҰҒА академигі және вице-президенті

Р е д а к ц и я а л қ а с ы:

Байзақов С.Б., э.ғ.д, проф., ҚР ҰҒА академигі (бас редактордың орынбасары); **Тиреуов К.М.**, э.ғ.д, проф., ҚР ҰҒА академигі (бас редактордың орынбасары); **Елешев Р.Е.**, т.ғ.д, проф., ҚР ҰҒА академигі; **Рау А.Г.**, т.ғ.д, проф., ҚР ҰҒА академигі; **Иванов Н.П.**, в.ғ.д, проф., ҚР ҰҒА академигі; **Кешуов С.А.**, т.ғ.д, проф., ҚР ҰҒА академигі; **Мелдебеков А.**, а.ш.ғ.д., проф., ҚР ҰҒА академигі; **Чоманов У.Ч.**, т.ғ.д, проф., ҚР ҰҒА академигі; **Елюбаев С.З.**, а.ш.ғ.д., проф., ҚР ҰҒА академигі; **Садыкулов Т.**, а.ш.ғ.д., проф., академигі; **Баймұқанов Д.А.**, а.ш.ғ.д., проф., ҚР ҰҒА корр-мүшесі; **Сансызбай А.Р.**, а.ш.ғ.д., проф., ҚР ҰҒА корр-мүшесі; **Умбетаев И.**, а.ш.ғ.д., проф., ҚР ҰҒА академигі; **Оспанов С.Р.**, а.ш.ғ.д., проф., ҚР ҰҒА құрметті мүшесі; **Олейченко С.И.**, а.ш.ғ.д., проф.; **Кененбаев С.Б.**, а.ш.ғ.д., проф., ҚР ҰҒА корр-мүшесі; **Омбаев А.М.**, а.ш.ғ.д., проф. ҚР ҰҒА корр-мүшесі; **Молдашев А.Б.**, э.ғ.д, проф., ҚР ҰҒА құрметті мүшесі; **Сагитов А.О.**, б.ғ.д, ҚР ҰҒА академигі; **Сапаров А.С.**, а.ш.ғ.д., проф., ҚР АШҒА академигі; **Балгабаев Н.Н.**, а.ш.ғ.д., проф.; **Умирзаков С.И.**, т.ғ.д, проф.; **Султанов А.А.**, в.ғ.д, проф., ҚР АШҒА академигі; **Алимкулов Ж.С.**, т.ғ.д, проф., ҚР АШҒА академигі; **Сарсембаева Н.Б.**, в.ғ.д, проф.

Р е д а к ц и я к е ñ е с і:

Fasler-Kan Elizaveta, Dr., University of Basel Switzerland; **Koolmees Petrus Adrianus**, Prof. Dr., Utrecht University, The Netherlands; **Babadoost-Kondri Mohammad**, Prof., University of Illinois, USA; **Yus Aniza Binti Yusof**, Dr., University Putra, Malaysia; **Hesseln Hayley Fawn**, As. Prof., University of Saskatchewan, Canada; **Alex Morgounov**, Pr., International Maize and Wheat Improvement Center Turkey; **Андреш С.**, Молдова Республикасы ҰҒА академигі; **Гаврилюк Н.Н.**, Украина ҰҒА академигі; **Герасимович Л.С.**, Беларусь Республикасының ҰҒА академигі; **Мамедов Г.**, Азербайжан Республикасының ҰҒА академигі; **Шейко И.П.**, Беларусь Республикасының ҰҒА академигі; **Жалнин Э.В.**, т.ғ.д., проф., Ресей; **Боинчан Б.**, а.ш.ғ.д, проф., Молдова Республикасы; **Юлдашбаев Ю.А.**, а.ш.ғ.д, проф., РФА корр-мүшесі, Ресей.

Главный редактор

Есполов Т.И.,

доктор эконом. наук, проф.,
вице-президент и академик НАН РК

Редакционная коллегия:

Байзаков С.Б., доктор эконом. наук, проф., академик НАН РК (заместитель главного редактора); **Тиреуов К.М.**, доктор эконом. наук, проф., академик НАН РК (заместитель главного редактора); **Елешев Р.Е.**, доктор техн. наук, проф., академик НАН РК; **Рау А.Г.**, доктор техн. наук, проф., академик НАН РК; **Иванов Н.П.**, доктор ветеринар. наук, проф., академик НАН РК; **Кешуов С.А.**, доктор техн. наук, проф., академик НАН РК; **Мелдебеков А.**, доктор сельхоз. наук, проф., академик НАН РК; **Чоманов У.Ч.**, доктор техн. наук, проф., академик НАН РК; **Елюбаев С.З.**, доктор сельхоз. наук, проф., академик НАН РК; **Садыкулов Т.**, доктор сельхоз. наук, проф., академик НАН РК; **Баймуқанов Д.А.**, доктор сельхоз. наук, проф., член-корр. НАН РК; **Сансызбай А.Р.**, доктор сельхоз. наук, проф., член-корр. НАН РК; **Умбетаев И.**, доктор сельхоз. наук, проф., академик НАН РК; **Оспанов С.Р.**, доктор сельхоз. наук, проф., Почетный член НАН РК; **Олейченко С.И.**, доктор сельхоз. наук, проф.; **Кененбаев С.Б.**, доктор сельхоз. наук, проф., член-корр. НАН РК; **Омбаев А.М.**, доктор сельхоз. наук, проф. член-корр. НАН РК.; **Молдашев А.Б.**, доктор эконом. наук, проф., Почетный член НАН РК; **Сагитов А.О.**, доктор биол. наук, академик НАН РК; **Сапаров А.С.**, доктор сельхоз. наук, проф., академик АСХН РК; **Балгабаев Н.Н.**, доктор сельхоз. наук, проф.; **Умирзаков С.И.**, доктор техн. наук, проф.; **Султанов А.А.**, доктор ветеринар. наук, проф., академик АСХН РК; **Алимкулов Ж.С.**, доктор техн. наук, проф., академик АСХН РК; **Сарсембаева Н.Б.**, доктор ветеринар. наук, проф.

Редакционный совет:

Fasler-Kan Elizaveta, Dr., University of asel Switzeland; **Koolmees Petrus Adrianus**, Prof. Dr., Utrecht University, The Netherlands; **Babadoost-Kondri Mohammad**, Prof., University of Illinois, USA; **Yus Aniza Binti Yusof**, Dr., University Putra, Malaysia; **Hesseln Hayley Fawn**, As.Prof., University of Saskatchewan, Canada; **Alex Morgounov**, Pr., International Maize and Wheat Improvement Center Turkey; **Андреш С.**, академик НАН Республики Молдова; **Гаврилюк Н.Н.**, академик НАН Украины; **Герасимович Л.С.**, академик НАН Республики Беларусь; **Мамедов Г.**, академик НАН Республики Азербайджан; **Шейко И.П.**, академик НАН Республики Беларусь; **Жалнин Э.В.**, доктор техн. наук, проф., Россия; **Боинчан Б.**, доктор сельхоз. наук, проф., Республика Молдова; **Юлдашбаев Ю.А.**, доктор сельхоз. наук, проф., член-корр. РАН, Россия.

Известия Национальной академии наук Республики Казахстан. Серия аграрных наук.

ISSN 2224-526X

Собственник: ООО «Национальная академия наук Республики Казахстан» (г. Алматы)

Свидетельство о постановке на учет периодического печатного издания в Комитете информации и архивов Министерства культуры и информации Республики Казахстан № 10895-Ж, выданное 30.04.2010 г.

Периодичность 6 раз в год

Тираж: 300 экземпляров

Адрес редакции: 050010, г. Алматы, ул. Шевченко, 28, ком. 219-220, тел. 272-13-19, 272-13-18

<http://nauka-nanrk.kz/agricultural.kz>

© Национальная академия наук Республики Казахстан, 2018

Адрес типографии: ИП «Аруна», г. Алматы, ул. Муратбаева, 75

Chief Editor

Espolov T.I.,

Dr. economy. Sciences, prof.,
Vice President and academician of the NAS RK

Editorial Board:

Baizakov S.B., Dr. of economy sciences, prof., academician of NAS RK (deputy editor); **Tireuov K.M.**, Doctor of Economy Sciences., prof., academician of NAS RK (deputy editor); **Eleshev R.E.**, Dr. Of agricultural sciences, prof., academician of NAS RK; **Rau A.G.**, Dr. sciences, prof., academician of NAS RK; **Ivanov N.P.**, Dr. of veterinary sciences, prof., academician of NAS RK; **Keshuov S.A.**, Dr. sciences, prof., academician of NAS RK; **Meldebekov A.**, doctor of agricultural sciences, prof., academician of NAS RK; **Chomanov U.Ch.**, Dr. sciences, prof., academician of NAS RK; **Yelyubayev S.Z.**, Dr. of agricultural sciences, prof., academician of NAS RK; **Sadykulov T.**, Dr. Farm. Sciences, prof., academician of NAS RK; **Baimukanov D.A.**, doctor of agricultural sciences, prof., corresponding member NAS RK; **Sansyzbai A.R.**, doctor of agricultural sciences, prof., corresponding member NAS RK; **Umbetaev I.**, Dr. Farm. Sciences, prof., academician of NAS RK; **Ospanov S.R.**, Dr. agricultural sciences, prof., Honorary Member of NAS RK; **Oleychenko S.N.**, Dr. Of agricultural sciences, prof.; **Kenenbayev S.B.**, Dr. Agricultural sciences, prof., corresponding member NAS RK; **Ombayev A.M.**, Dr. Agricultural sciences, Prof. corresponding member NAS RK; **Moldashev A.B.**, Doctor of Economy sciences, prof., Honorary Member of NAS RK; **Sagitov A.O.**, Dr. biol. sciences, academician of NAS RK; **Saparov A.S.**, Doctor of agricultural sciences, prof., academician of NAS RK; **Balgabaev N.N.**, the doctor agricultural sciences, Prof.; **Umirzakov S.I.**, Dr. Sci. Sciences, Prof.; **Sultanov A.A.**, Dr. of veterinary sciences, prof., academician of the Academy of Agricultural Sciences of Kazakhstan; **Alimkulov J.C.**, Dr. of tekhncial sciences, prof., academician of the Academy of Agricultural sciences of Kazakhstan; **Sarsembayeva N.B.**, Dr. veterinary sciences, prof.

Editorial Board:

Fasler-Kan Elizaveta, Dr., University of Basel Switzzeland; **Koolmees Petrus Adrianus**, Prof. Dr., Utrecht University, The Netherlands; **Babadoost-Kondri Mohammad**, Prof., University of Illinois, USA; **Yus Aniza Binti Yusof**, Dr., University Putra, Malaysia; **Hesseln Hayley Fawn**, As. Prof., University of Saskatchewan, Canada; **Alex Morgounov**, candidate of agricultural sciences, International Maize and Wheat Improvement Center Turkey; **Andresh S.**, academician of NAS of Moldova; **Gavriluk N.N.**, academician of NAS of Ukraine; **Gerasimovich L.S.**, academician of NAS of Belorassia; **Mamadov G.**, academician of NAS of Azerbaijan; **Sheiko I.P.**, academician of NAS of Belorassia; **Zhalnin E.V.**, Dr. of technical sciences, professor, Russia, **Boinchan B.**, doctor of agricultural sciences, prof., Moldova; **Yuldashbayev Y.A.**, doctor of agricultural sciences, prof., corresponding member of RAS, Russia.

News of the National Academy of Sciences of the Republic of Kazakhstan. Series of Agrarian Sciences.

ISSN 2224-526X

Owner: RPA "National Academy of Sciences of the Republic of Kazakhstan" (Almaty)

The certificate of registration of a periodic printed publication in the Committee of Information and Archives of the Ministry of Culture and Information of the Republic of Kazakhstan N 10895-Ж, issued 30.04.2010

Periodicity: 6 times a year

Circulation: 300 copies

Editorial address: 28, Shevchenko str., of.219-220, Almaty, 050010, tel. 272-13-19, 272-13-18,
<http://nauka-nanrk.kz/agricultural.kz>

© National Academy of Sciences of the Republic of Kazakhstan, 2018

Address of printing house: ST "Aruna", 75, Muratbayev str, Almaty

NEWS

OF THE NATIONAL ACADEMY OF SCIENCES OF THE REPUBLIC OF KAZAKHSTAN

SERIES OF AGRICULTURAL SCIENCES

ISSN 2224-526X

<https://doi.org/10.32014/2018.2224-526X.2>

Volume 5, Number 47 (2018), 12 – 19

K. Zh. Tagayev¹, A. I. Morgounov², M. A. Yessimbekova³,
A. I. Abugalieva³, G. O. Bayadilova¹

¹Kazakh National Agrarian University, Almaty, Kazakhstan,

²International Maize and Wheat Improvement Center (CIMMYT), Ankara, Turkey,

³Kazakh Research Institute of Agriculture and Plant Growing, Almalyk village, Almaty region, Kazakhstan

EFFECT OF COMMON BUNT INFECTION ON AGRONOMIC TRAITS AND RESISTANCE IN WINTER WHEAT LINES

Abstract. Common bunt (*Tilletia caries*) is a seed transmitted fungal disease in wheat. The resistance cultivars and lines of wheat should use to control this type of diseases in organic farming. A set of 75 wheat cultivars and lines from IWWIP of Turkey used. During the period 2016-2017, an experiment was carried out at the Kazakh Research Institute of Agriculture and Growing in an artificially inoculated nursery. The susceptible check, GEREK 79, had a high level of susceptibility to common bunt with 59.7% infected heads. The high mean disease incidence in the nursery was 74.4%. The sixteen genotypes were resistant to disease under artificial inoculation. The forty-two wheat genotypes (56% of all genotypes) expressed moderate resistance, which infected around 2.0-27.3% of ears. The wheat lines had different levels of agronomic traits under artificial inoculation of common bunt. The productivity of wheat genotypes under artificial infection ranged from 1.13 to 7.29 t/ha. The expected strong positive correlation was detected between the grain number and grain weight ($r = 0.7$), between the grain weight and TKW ($r = 0.75$) and the grain weight and total grain weight ($r = 0.79$). The negative correlation was found between the bunted ears and all agronomic traits. Identified resistance genotypes will be useful for breeding programs to forming resistance cultivars to common bunt in Kazakhstan.

Keywords: wheat, wheat cultivars, wheat lines, common bunt, resistance, productivity.

The Kazakhstan is one of the major wheat producers in the world. The climatic conditions on the north are very favorable to cultivate cereals crops. The total area planted to wheat in the country represents over 85% of total cereal production. Currently Kazakhstan produces 18-20 million tons of wheat grain, but output is highly dependent on weather and in recent years has fluctuated between 10 and 17 million tons. Kazakhstan is a major exporter of wheat and plays an important role in the food security of Central Asia [1]. However, diseases and pests also play an important role in yield reduction in Kazakhstan. Common bunt, caused by *Tilletiatritici* (Bjerk.) G. Wint. *In* Rabenh. [syn. *T. caries* (DC.) Tul. & C. Tul.) and *T. laevis* Kuhn *in* Rabenh. (syn. *T. foetida* (Wallr.) Liro.), has occurred in all wheat-growing countries of the world [2, 3]. The disease is largely spread in all wheat-growing regions [4]. The spores which are left on the surface of the contaminated seeds are systematically developing and multiplying, year by year, inside the growing wheat plants, when the plant reaches maturity, they transform the core of the wheat kernels into toxic fungus spores. In case of heavy incidence it is not possible to use the seed as food or feed. It causes yield loss in common wheat [5, 6] and reduces grain quality [7]. Already low doses of the spores represent a problem for seed sales and multiplication. The spores contain trimethylamine causing an unpleasant odour. Although chemical seed treatments can effectively control these diseases, especially common bunt, resistant cultivars remain desirable for bunt management in developing countries [8], for organic production of wheat, and as a lower-cost alternative to chemical treatment [9]. Although the chemical treatment of wheat seed for the control of common bunt is widely used, genetic resistance of wheat is an important part of the bunt control in many countries, especially on organic farms. Resistant varieties may reduce the losses due to bunt infection [10]. The most economic and effective way of

controlling common bunt of wheat is using bunt resistance cultivars [11-13]. It is important to identify sources of wheat resistance to common bunt in order to use resistant varieties and to develop resistant cultivars through breeding effort. Testing for resistance to *Tilletia spp.* and identification of new sources of resistance are necessary especially under organic farming conditions [14]. Estimating the level of common bunt resistance is time consuming due to the need to wait until the symptoms are expressed. Symptoms happen when the grain filling period is nearly complete. The symptoms sometimes are only expressed on the last spikes formed, and the symptoms often are only expressed in a few of the florets [11]. So, understanding the effect of the common bunt on earlier agronomic traits would help in identifying the trait associated with the resistance and hence the ability to select resistant genotypes earlier rather than waiting until plant maturity. For example, it was reported that common bunt has an effect on plant height, number of heads and root length of heads [15].

Our aim of this study was to evaluate agronomic traits and resistance of wheat cultivars and lines to common bunt under artificial infection. The approach of this study was to study the effect of common bunt infection on some easily measured agronomic traits in a set of seventy-five International common bunt nursery's winter wheat genotypes. Also identification resistant and susceptible genotypes to know if there are differences in the response of the agronomic traits to the disease in order to determine if we can use an agronomic trait as an indicator for the level of resistance to common bunt.

Materials and methods. A set of 75 winter wheat lines from Common Bunt Resistant Nursery (CBUNT- RN 2015-2016) were used to study effect of common bunt on some important agronomic traits. This nursery was performed and distributed by the International Winter Wheat Improvement Program of Turkey and combined different type of germplasm resistant to common bunt (75 wheat genotypes from 7 countries –Turkey, Iran, Kazakhstan, Mexico, Romania, Russia, and USA). The experiments of artificial inoculation with spores of common bunt were organized in the field, in the agricultural research blocks of Kazakh Research Institute of Agriculture and Growing, Almalybak, Almaty region, Kazakhstan, in the period 2016-2017. The experiments were carried out in randomized complete blocks with three repetitions. After inoculation, approximately 150 seeds have sown from each germplasm. Artificial inoculation of the seeds with spores has done by methods A.I. Borrgardt-Anpilogova [16], using a mixture of isolates of southeast part of Kazakhstan. The spores were obtained sampling natural infected plants from the field, from Scientific Research Institute for Biological Safety Problems, Otar, Kazakhstan. The doses of spores used for inoculation vary a lot [17]. In order to provide a total infection, the wheat cultivars and lines were treated separately with a higher inoculation dose of 0.08 g spores/150 seeds [18]. A winter wheat resistant (MUFITBEY and NACIBEY) and susceptible checks (GEREK79) were included to confirm the effectiveness of the common bunt inoculation protocol. The seeds were sown in the autumn and in the next year, at the heading stage, the disease could be detected in the field. The specific symptoms were evaluated at the heading stage by the visual inspection of the spikes in the month of June, when we could observe, at the infected plants, the dark coloured spikes, the spreading of the glumes and, at the maturity, the grains content transformed into a mass of dark coloured spores. The evaluation tests provided by scale Krivchenko, level of percent (%) of infected ears [19]. The resistance evaluation have done by reporting the number of susceptible plants to the total number of plants. The plant height was measured during ripening stage as the height of the plant from the ground to the tip of the plant. The length of the head was measured from the base of the head to the tip after the plants completely emerged. Number of spike was measured by counting of all spikes in one plant. The number of grain per spike was measured by counting all the grains in one general spike. The grain weight per spike, 1000 kernel weight and total grain weight were measured on mature plants that had not been watered for seven days as the average grain weight, weight of 1000 grains and weight of all grains, respectively. Statistical and correlation analysis were provided using Excel 2010.

Results and discussion. Resistance to common bunt. A mixture of common bunt teliospores from races for field in southeast part of Kazakhstan induced a different reaction on wheat lines of International CBUNT Nursery. The winter susceptible check, GEREK 79, had a high level of susceptibility to common bunt with 59.7% infected heads (table). This high level of infection in the susceptible check on cultivar GEREK 79 confirmed that the common bunt infection was successful. Goates (1996) [3] suggested that common bunt resistance evaluation should be considered valid when a susceptible check had more than 50% infected heads. The two resistance winter genotypes, MUFITBEY and NACIBEY, had similar

degrees of resistance with 10.1 and 2.0% infected heads, respectively. The high mean disease incidence in the nursery was 74.4%. Among 75 lines tested genotypes from CBUNT International nursery, 16 genotypes were resistant to disease under artificial inoculation:

PBW343*2/KUKUNA//ATAY/CALVEZ/3/ATAY/GALVEZ87, ORKINOS-1/SUNR23//SONMEZ, ATAY/GALVEZ87/6/TAST/SPRW/4/ROM-TAST/BON/3/DIDO//SU92/CI13645/5/F130L.12, MADSEN/MALCOLM//ZARGANA-9/3/BURBOT-6, RINA-6/ORKINOS-7, DE9//MERGAN-2, ORKINOS-1*2/3/AUS GS50AT34/SUNCO//CUNNINGHAM, KS902709-B-5-1/BURBOT-4, RANA96/GANSU-3, RINA-6/BEZ/NAD//KZM(ES85.24)/3/F900K, ALMT*3/7/VEE/CMH77A.917//VEE/6/CMH79A.955/4/AGA/3/SN64*4/CNO67//INIA66/5/NAC, BEZOSTAYA/AE.CYLINDRICA, BEZOSTAYA/TR.MILITINAE//TR.MILITINAE-6, BEZOSTAYA/TR.MILITINAE// TR.MILITINAE-4, CV.RODINA/AE/SPELTOIDES(10 KR) and OSTROV. It is 21.3% of all studied wheat genotypes. The other 42 wheat genotypes (56% of all genotypes) expressed moderate resistance, which infected around 2.0-27.3% of ears. Ten wheat lines were susceptible (31.4-48.2% infected ears) and seven lines were very susceptible (up to 50% infected ears) to common bunt infection.

Common bunt resistance and agronomic traits of wheat genotypes

| # | Name of wheat lines | Plant height, Cm | Head length, Cm | Spike number | Grain number/ Spike | Grain weight/ spike, g | 1000 kernel weight, g | Total grain weight, g | % bunte-dears | Grain yield, t/ha |
|----|--|------------------|-----------------|--------------|---------------------|------------------------|-----------------------|-----------------------|---------------|-------------------|
| 1 | 2 | 3 | 4 | 5 | 6 | 7 | 8 | 9 | 10 | 11 |
| 1 | MUFITBEY (resistant check) | 120,0 | 12,0 | 25,0 | 83,0 | 3,8 | 45,7 | 30,5 | 10,1 | 6,08 |
| 2 | NACIBEY (resistant check) | 105,0 | 10,0 | 19,0 | 61,0 | 1,8 | 29,5 | 16,6 | 2,0 | 2,17 |
| 3 | GEREK79 (susceptible check) | 116,0 | 9,0 | 19,0 | 47,0 | 0,7 | 14,8 | 12,7 | 39,7 | 2,63 |
| 4 | PBW343*2/KUKUNA//ATAY/CALVEZ/3/ATAY/GALVEZ87 | 85,0 | 11,0 | 21,0 | 56,0 | 1,0 | 17,8 | 14,3 | 0,0 | 3,68 |
| 5 | 87-461 a 63-555//SAULESKU#26/PARUS/3/AGRI/NAC//ATTILA | 85,0 | 11,0 | 25,0 | 80,0 | 2,7 | 33,7 | 18,5 | 57,2 | 4,44 |
| 6 | ORKINOS-1/SUNR23//SONMEZ | 115,0 | 13,0 | 23,0 | 79,0 | 3,5 | 44,3 | 22,2 | 0,0 | 4,50 |
| 7 | ATAY/GALVEZ87/6/TAST/SPRW/4/ROM-TAST/BON/3/DIDO//SU92/CI13645/5/F130L.12 | 125,0 | 14,0 | 23,0 | 79,0 | 3,7 | 46,8 | 28,7 | 0,0 | 3,35 |
| 8 | MADSEN/MALCOLM//ZARGANA-9/3/BURBOT-6 | 130,0 | 12,0 | 21,0 | 52,0 | 2,0 | 38,4 | 12,5 | 0,0 | 4,17 |
| 9 | RINA-6/ORKINOS-7 | 115,0 | 12,0 | 25,0 | 64,0 | 2,4 | 37,5 | 15,0 | 0,0 | 2,58 |
| 10 | SAULESKU#44/TR810200//GRISSET-4 | 100,1 | 12,0 | 21,0 | 73,0 | 2,2 | 30,1 | 20,0 | 57,2 | 7,29 |
| 11 | ATTILA/BABAX//PASTOR/4/TAST/SPRW//ZAR/3/ATAY/GALVEZ87 | 90,1 | 12,0 | 23,0 | 78,0 | 2,9 | 37,1 | 22,0 | 47,8 | 2,77 |
| 12 | BURBOT-4/3/OMBUL/ALAMO//MV11 | 100,1 | 12,0 | 23,0 | 71,0 | 2,4 | 33,8 | 19,0 | 6,1 | 2,12 |
| 13 | FRTL//AGRI//NAC/3/BONITO-36/4/ERIT58-87//KS82W409/SPN/3/KRC66/SERI | 105,1 | 10,0 | 19,0 | 50,0 | 1,7 | 34,0 | 10,2 | 11,9 | 1,92 |
| 14 | GUN91/MNCH*2//T-2003 | 105,1 | 11,0 | 21,0 | 67,0 | 2,6 | 39,0 | 21,0 | 15,4 | 3,63 |
| 15 | KRASnodAR/FRTL/6/NGDA146/4/YMN/TOB//MCD/3/LIRA/5/F130L.12 | 120,1 | 11,0 | 25,0 | 84,0 | 3,5 | 41,6 | 25,6 | 11,1 | 6,24 |
| 16 | TJB368-251/BUC//SMUT1590-165/3/KS7866-15/ORS8425/4/NE87U119/CHAM//1D13.1/MKT | 125,1 | 13,0 | 23,0 | 71,0 | 3,2 | 45,0 | 24,5 | 4,6 | 1,80 |
| 17 | SHARK/F44105W2.1//AUS4930.7/2*PA STOR/3/ORKINOS-1 | 100,1 | 10,0 | 21,0 | 78,0 | 2,3 | 29,4 | 19,1 | 26,5 | 6,08 |
| 18 | GANSU-1/3/AUS GS50AT34/SUNCO//CUNNINGHAM/4/ORKINOS-1 | 95,1 | 12,0 | 21,0 | 63,0 | 2,8 | 44,4 | 27,2 | 4,2 | 4,83 |
| 19 | BURBOT-4/3/OMBUL/ALAMO//MV11 | 95,1 | 11,0 | 21,0 | 60,0 | 1,5 | 25,0 | 17,0 | 9,3 | 4,81 |
| 20 | KUPAVA/BURBOT-4//PYN/2*BAU | 90,2 | 12,0 | 21,0 | 57,0 | 2,2 | 38,5 | 16,0 | 54,8 | 3,05 |
| 21 | DE9//MERGAN-2 | 125,2 | 14,0 | 21,0 | 59,0 | 2,5 | 42,3 | 21,0 | 0,0 | 2,12 |

| <i>Continuation of table</i> | | | | | | | | | | |
|------------------------------|---|-------|------|------|------|-----|------|------|------|------|
| 1 | 2 | 3 | 4 | 5 | 6 | 7 | 8 | 9 | 10 | 11 |
| 22 | KRASNODAR/FRTL/6/NGDA146/4/YMN/TOB//MCD/3/LIRA/5/F130L.12 | 125,2 | 12,0 | 23,0 | 70,0 | 2,8 | 40,0 | 21,8 | 8,6 | 6,17 |
| 23 | 362K2.111//TX71A1039.VI*3/AMI/3/ES14/130L.12//MNCH | 120,2 | 11,0 | 21,0 | 74,0 | 2,0 | 27,0 | 18,6 | 10,0 | 5,20 |
| 24 | SELYNKA/MERGAN-1 | 125,2 | 10,0 | 19,0 | 50,0 | 1,8 | 36,0 | 15,2 | 23,1 | 1,70 |
| 25 | 91-142 A 61/KATIA1//GRIZET-4 | 120,2 | 11,0 | 21,0 | 65,0 | 2,6 | 40,0 | 18,2 | 10,0 | 4,70 |
| 26 | KUPAVA/BURBOT-4//PYN/2*BAU | 100,2 | 11,0 | 19,0 | 50,0 | 1,3 | 26,0 | 16,5 | 31,4 | 5,67 |
| 27 | KRASNODAR/FRTL/6/NGDA146/4/YMN/TOB//MCD/3/LIRA/5/F130L.12 | 115,2 | 11,0 | 23,0 | 65,0 | 2,7 | 41,5 | 16,0 | 34,5 | 4,25 |
| 28 | ORKINOS-1*2/3/AUSGS50AT34/SUNCO//CUNNINGHAM | 135,2 | 12,0 | 23,0 | 57,0 | 2,4 | 42,1 | 19,2 | 0,0 | 4,35 |
| 29 | 87-461 a 63-555/4/ERIT58-87//KS82W409//SPN/3/KRC66/SERI | 110,2 | 13,0 | 23,0 | 57,0 | 2,5 | 43,8 | 19,3 | 32,5 | 1,55 |
| 30 | SAULESKU#44/TR810200//ZGI | 115,3 | 12,0 | 21,0 | 57,0 | 2,1 | 36,8 | 19,5 | 50,0 | 4,50 |
| 31 | KRASNODAR/FRTL/6/NGDA146/4/YMN/TOB//MCD/3/LIRA/5/F130L.12 | 115,3 | 12,0 | 25,0 | 63,0 | 2,7 | 42,8 | 21,5 | 15,4 | 5,50 |
| 32 | TAM200/KAUZ/4/CHAM6//1D13.1/MLT/3/SHI4414/CROW | 80,3 | 13,0 | 23,0 | 80,0 | 3,3 | 41,2 | 18,8 | 21,2 | 2,80 |
| 33 | SHARK/F44105W2.1//CHARA/3/MERGAN-1 | 100,3 | 10,0 | 21,0 | 65,0 | 2,4 | 37,0 | 17,6 | 15,6 | 3,98 |
| 34 | ALPU/VR5053(WA#FM/201/23*2/GS50A) | 105,3 | 12,0 | 23,0 | 62,0 | 2,0 | 32,2 | 16,4 | 25,0 | 3,73 |
| 35 | KS902709-B-5-1/BURBOT-4 | 110,3 | 11,0 | 21,0 | 55,0 | 2,1 | 38,1 | 18,8 | 0,0 | 5,55 |
| 36 | JCAM/EMU//DOVE/3/JGR/4/THK/5/BOEMA | 95,3 | 11,0 | 21,0 | 80,0 | 2,2 | 27,5 | 18,0 | 12,9 | 3,20 |
| 37 | BATERA//KEA/TOW/3/TAM200/4/494J6.11/TRAP#1/BOW/5/TX96V2427 | 105,3 | 10,0 | 21,0 | 55,0 | 1,6 | 29,0 | 12,8 | 6,1 | 2,70 |
| 38 | BATERA//KEA/TOW/3/TAM200/4/494J6.11/TRAP#1/BOW/5/TX96V2427 | 95,3 | 12,0 | 21,0 | 63,0 | 2,0 | 31,7 | 15,1 | 5,9 | 5,27 |
| 39 | ORKINOS-1/4/JING411//PLK70/LIRA/3/GUN91 | 125,3 | 14,0 | 25,0 | 76,0 | 3,1 | 40,7 | 25,8 | 15,4 | 2,12 |
| 40 | GRIZET-4/3/ID#840335//PIN39/PEW/4/LILIA BG/GT | 120,4 | 13,0 | 23,0 | 90,0 | 2,8 | 31,1 | 17,8 | 17,9 | 4,55 |
| 41 | KAMBARA1/ZANDER-17 | 95,4 | 14,0 | 25,0 | 72,0 | 2,1 | 29,1 | 18,7 | 27,3 | 2,20 |
| 42 | ADMIS/5/SMB/HN4//SPN/3/WTS//YMH/HYS/4/SAB | 100,4 | 11,0 | 21,0 | 66,0 | 2,3 | 34,8 | 20,8 | 17,7 | 6,45 |
| 43 | RANA96/GANSU-3 | 125,4 | 12,0 | 23,0 | 74,0 | 3,1 | 41,8 | 21,2 | 0,0 | 5,05 |
| 44 | RINA-6/BEZ/NAD//KZM(ES85.24)/3/F900K | 110,4 | 11,0 | 22,0 | 49,0 | 2,1 | 42,8 | 14,3 | 0,0 | 5,35 |
| 45 | VORONA/3/TOB*2/7C//BUC/4/CHAM6//1D13.1/MLT/3/SHI4414/CROW | 110,4 | 9,0 | 21,0 | 58,0 | 2,0 | 34,4 | 15,0 | 5,3 | 4,83 |
| 46 | Son64/4/Wr51/mida//Nt.h/K117/5/Anza/3/Pi//Nor/Hys/4/ Sefid | 120,4 | 11,0 | 19,0 | 58,0 | 2,8 | 48,2 | 23,4 | 20,0 | 5,24 |
| 47 | ALD/SNB//ZARRIN/3/YACO/2*PARUS | 70,4 | 9,0 | 19,0 | 72,0 | 2,8 | 38,8 | 18,7 | 33,3 | 1,93 |
| 48 | SPN/MCD//CAMA/3/NZR/4/ALD/SNB*2/5/GASCOGNE | 75,4 | 11,0 | 21,0 | 61,0 | 1,3 | 21,3 | 7,8 | 6,5 | 2,78 |
| 49 | SPN/MCD//CAMA/3/NZR/4/ALD/SNB*2/5/GASCOGNE | 65,4 | 10,0 | 20,0 | 55,0 | 1,3 | 23,6 | 11,5 | 5,9 | 2,58 |
| 50 | CMH79A.955/4/AGA/3/4*SN64/CNO67//INIA66/5/NAC/6/CMH83.25//RSH/8/ZRN | 80,5 | 11,0 | 23,0 | 72,0 | 2,2 | 27,7 | 18,1 | 21,7 | 2,85 |
| 51 | CMH79A.955/4/AGA/3/4*SN64/CNO67//INIA66/5/NAC/6/CMH83.25//RSH/8/ZRN | 90,5 | 12,0 | 23,0 | 83,0 | 2,5 | 30,1 | 18,0 | 13,9 | 2,78 |
| 52 | CMH79A.955/4/AGA/3/4*SN64/CNO67//INIA66/5/NAC/6/CMH83.25//RSH/8/ZRN | 85,5 | 12,0 | 21,0 | 64,0 | 1,8 | 28,1 | 16,0 | 17,5 | 2,65 |

| <i>Continuation of table</i> | | | | | | | | | | |
|------------------------------|--|-------|------|------|------|-----|------|------|------|------|
| 1 | 2 | 3 | 4 | 5 | 6 | 7 | 8 | 9 | 10 | 11 |
| 53 | QUDS*3/MV17 | 95,5 | 11,0 | 21,0 | 67,0 | 2,0 | 28,3 | 16,2 | 48,2 | 2,63 |
| 54 | ALMT*3/7/VEE/CMH77A.917//VEE/6/CMH79A.955/4/AGA/3/SN64*4/CNO67//INIA66/5/NAC | 80,6 | 10,0 | 21,0 | 53,0 | 1,8 | 34,0 | 14,0 | 0,0 | 4,83 |
| 55 | CROC_1/AE.SQUARROSA(224)/OPAT A | 80,5 | 12,0 | 19,0 | 52,0 | 1,8 | 35,7 | 14,7 | 20,6 | 3,50 |
| 56 | SANZAR-8/KKTS | 90,5 | 11,0 | 21,0 | 72,0 | 2,0 | 30,5 | 16,7 | 41,9 | 3,68 |
| 57 | INTENSIVNAYA//PBW343*2//TUKURU | 90,5 | 12,0 | 19,0 | 64,0 | 2,0 | 29,6 | 14,5 | 23,4 | 2,83 |
| 58 | TSAPKI/FARMEC | 90,5 | 11,0 | 21,0 | 71,0 | 2,0 | 26,7 | 20,1 | 57,2 | 6,47 |
| 59 | AMCEL/KS970274/3/KS91048L-2-1/CM112793(CHL)/2*STAR)/HWK1064-6 | 95,5 | 12,0 | 19,0 | 56,0 | 2,3 | 41,0 | 19,2 | 2,2 | 2,53 |
| 60 | DORADE-5/KS980512 | 100,6 | 12,0 | 23,0 | 92,0 | 3,5 | 38,0 | 21,0 | 23,8 | 6,35 |
| 61 | OR 943576/KS920709 | 95,6 | 13,0 | 23,0 | 85,0 | 1,8 | 21,1 | 17,0 | 5,9 | 3,47 |
| 62 | MRS/CI14482//YMH/HYS/3/RONDEZVOUS/4/ABI 86*3414X84W063-9939-2//KARL92 | 90,6 | 13,0 | 21,0 | 60,0 | 1,8 | 30,0 | 20,0 | 44,5 | 4,15 |
| 63 | KS92WGRC-25 | 110,6 | 11,0 | 19,0 | 49,0 | 1,5 | 30,6 | 13,0 | 38,5 | 4,73 |
| 64 | BEZOSTAYA/AE.CYLINDRICA | 120,6 | 14,0 | 29,0 | 81,0 | 3,0 | 37,0 | 25,3 | 0,0 | 4,51 |
| 65 | BEZOSTAYA/TR.MILITINAE//TR.MILITINAE-6 | 140,6 | 16,0 | 24,0 | 97,0 | 3,0 | 30,0 | 20,5 | 0,0 | 1,55 |
| 66 | BEZOSTAYA/TR.MILITINAE//TR.MILITINAE-4 | 140,5 | 14,0 | 24,0 | 49,0 | 1,8 | 36,7 | 20,7 | 0,0 | 3,68 |
| 67 | CV.RODINA/AE/SPELTOIDES(10 KR)/S.CEREALE(1.OKR) | 140,4 | 14,0 | 24,0 | 62,0 | 1,8 | 29,0 | 16,8 | 20,0 | 3,60 |
| 68 | CV.RODINA/AE/SPELTOIDES(10 KR)/S.CEREALE(1.OKR) | 140,0 | 14,0 | 24,0 | 59,0 | 2,0 | 33,8 | 18,7 | 5,4 | 4,37 |
| 69 | CV.RODINA/AE/SPELTOIDES(10 KR) | 140,7 | 14,0 | 24,0 | 71,0 | 2,6 | 36,6 | 20,2 | 0,0 | 5,68 |
| 70 | F06393GP10 | 70,7 | 10,0 | 22,0 | 63,0 | 1,8 | 28,5 | 16,7 | 25,8 | 0,87 |
| 71 | F08034G1 | 75,7 | 10,0 | 22,0 | 62,0 | 2,2 | 35,4 | 17,3 | 34,5 | 4,43 |
| 72 | F08347G8 | 75,8 | 11,0 | 22,0 | 74,0 | 3,0 | 39,1 | 19,0 | 8,9 | 1,13 |
| 73 | OSTROV | 75,7 | 11,0 | 20,0 | 79,0 | 2,7 | 34,1 | 17,0 | 0,0 | 2,32 |
| 74 | F07270G2 | 100,9 | 10,0 | 18,0 | 44,0 | 1,8 | 41,0 | 16,3 | 74,4 | 4,17 |
| 75 | F00628G34-1 | 80,7 | 10,0 | 18,0 | 80,0 | 3,1 | 38,7 | 24,1 | 4,9 | 3,53 |

Agronomic evaluation. In the present work 9 traits were assessed on 75 winter wheat lines through 2 years grown under artificial infection of common bunt. The wheat lines had different levels of agronomic traits under artificial inoculation of common bunt. The plant and spike height were 70.4-140.7 cm and 9.0-14.0 cm, respectively. The number of spike per plant ranged from 18 to 25 spikes. The grain weight per spike was 0.7-3.8 gram. The 1000 kernel weight and total grain weight were showed different level of productivity, 14.8-48.2 gram and 7.8-30.5 gram, respectively. The productivity of wheat genotypes under artificial infection ranged from 1.13 to 7.29 t/ha.

Correlation analysis. The correlations among the nine traits were mostly similar, but in some cases, the strength of the correlations was different. The correlation coefficients were found to be highly significant at the 0.05 probability level. The moderately positive correlation was detected between the plant height and head length ($r = 0.51$), between the head length and number of spike ($r = 0.62$). Also, moderately positive correlation was found between grain number and total grain weight ($r = 0.53$). The expected strong positive correlation was detected between the grain number and grain weight ($r = 0.7$), between the grain weight and TKW ($r = 0.75$) (figures 1, 2).




Figure 1 – The correlation coefficient of the between grain number and grain weight per spike




Figure 2 – The correlation coefficient of the between grain weight per spike and TKW




Figure 3 – The correlation coefficient of the between grain weight per spike and total grain weight per plant

Also, strong positive correlation was found between grain weight and total grain weight ($r = 0.79$) (figure 3). The negative correlation was found between the bunted ears and all agronomic traits, but these correlations were not so high, from $r = 0.1$ to $r = 0.29$.

Conclusion. In conclusion, the common bunt infection was found to decrease the productivity and biological yield in the tested genotypes. Artificial inoculation tests for common bunt resistance showed that a large number of resistance genotypes is available in disease condition of southeast part of Kazakhstan. On base this study sixteen genotypes from the CBUNT Nursery of IWWIP can be considered as valuable resistance sources to common bunt. Identified resistance genotypes will be useful for breeding programs to forming resistance cultivars to common bunt in Kazakhstan.

REFERENCES

- [1] Kokhmetova A., Sapakhova Z., Urazaliev R., Yessimbekova M., Yeleshev R., Morgounov A. Effect of spring biomass removal on expression of agronomic traits of winter wheat // World Applied Sciences Journal. 2014. N 30 (3). P. 322-329.
- [2] Munjal R.L. Bunt disease of wheat // Sci. Rep. 1966. N 3. P. 33-36.
- [3] Bahadur P., Singh B.M. Hill bunt of wheat and its importance // Int. J. Trop. Plant Dis. 1987. N 5. P. 25-33.
- [4] Wilcoxson R.D., Saari E.E. (eds.). Bunt and smut diseases of wheat: concepts and methods of disease management. Mexico, D.F.: CIMMYT, 1996. P. 66.
- [5] Holton C.S. Host selectivity as a factor in the establishment of physiologic races of *Tilletia caries* and *T. foetida* produced by hybridization // Phytopathology. 1947. N 37. P. 817-821.
- [6] Goel L.B., Singh D.V. Smuts and bunts of wheat and their control. In S. P. Raychaudhuri, A. Varma, K. S. Bhargava, and B. S. Mehrotra, eds. Advances in mycology and plant pathology. Harsh Kumar at Sagar Printers, New Delhi, India., 1975. P. 131-147.
- [7] Flor H.H., Gaines E.F., Smith W.K. The effect of bunt on yield of wheat // J. Am. Soc. Agron. 1932. N 24. P. 778-784.
- [8] Saari E.E., Mamluk O.F., Burnett P.A. Bunts and smuts of wheat. In R.D. Wilcoxson and E.E. Saari (ed.) Bunt and smut diseases of wheat: Concepts and methods of disease management. CIMMYT. Mexico, D.F., 1996. P. 1-11.
- [9] Bonman J.M., Bockelman H.E., Goates B.J., Obert D.E., McGuire P.E., Qualset C.O., Hijmans R.J. Geographic distribution of common and dwarf bunt resistance in landraces of *Triticum aestivum* subsp. *aestivum* // Crop Sci. 2006. N 46. P. 1622-1629.
- [10] Dumalasova V., Bartoš P. Resistance of winter wheat cultivars to common bunt, *Tilletiatriticici*(Bjerk.). Wint. and *T. laevis* Kühn. // Journal of Plant Diseases and Protection. 2006. N 113. P. 159-163.
- [11] Goates B.J. Common bunt and dwarf bunt. In: Wilcoxson R.D., Saari E.E. (eds.): Bunt and smut diseases of wheat: concepts and methods of disease management. Mexico, DF, CYMMIT, 1996. P. 12-25.
- [12] Smeltzer D.G. Inheritance of resistance to race T-1 of *Tilletia caries* in Minturki and Cooperatorka wheats // Agron. J. 1952. N 44. P. 529-533.
- [13] Gaudet D.A., Puchalski B.L. Status of bunt resistance in western Canadian spring wheat and triticale // Can. J. Plant Sci. 1989. N 69. P. 797-804.
- [14] Borgen A. Biological control of common bunt in organic agriculture // Journal of Crop Production. 2000. N 3. P. 159-174.
- [15] Mourad A., Mahdy E., Bakheit B.R., Abo-Elwafaa A., Baenziger P.S. Effect of common bunt infection on agronomic traits in wheat (*Triticum aestivum* L.) // J Plant Genet Breed. 2018. N 2(1). P. 102-109.
- [16] Anpilogova M.Z. Methods of resistance characteristics of parent materials to fungi diseases and select pair for crossing processes // Proc. Botanic, genetics and breeding. 1958. N 33(1). P. 200-320 (In Russ.).
- [17] Dumalasova V., Bartos P. Effect of inoculum doses on common bunt infection on wheat caused by *Tilletiatriticici* and *T. laevis* // Czech J. Genet. Plant. Breed. 2008. N 44(2). P. 73-77.
- [18] Nielsen B.J. Strategies for avoiding seed-borne diseases // Proceedings 11th Australian Barley Technical Symposium and 53rd Australian Cereal Chemistry Conference. 2003. P. 75-79.
- [19] Krivchenko V.I. Methods of evaluation resistance of cereal crops to bunt diseases. M.: Kolos, 1971. P. 304. (In Russ.).

К. Ж. Тағаев¹, А. И. Моргун², М. А. Есимбекова³, А. И. Аbugалиева³, Г. О. Баядилова¹

¹Қазақ ұлттық аграрлық университеті, Алматы, Қазақстан,

²Жүгері мен бидайды жақсарту халықаралық орталығы (СИММИТ), Анкара, Түркия,

³Қазақ егіншілік және өсімдік шаруашылығы ғылыми-зерттеу институты,
Алмалыбақ, Алматы облысы, Қазақстан

ҚАТТЫ ҚАРАКҮЙЕ АУРУЫНЫҢ КҮЗДІК БИДАЙ ЛИНИЯЛАРЫНЫҢ АГРОНОМИЯЛЫҚ БЕЛГІЛЕРІМЕН ТӨЗІМДІЛІГІНЕ ӘСЕРІ

Аннотация. Қатты қаракүйе (*Tilletia caries*) тұқым арқылы берілетін бидайдың саңырауқұлақ ауруларының бірі. Органикалық ауылшаруашылығында аурудың алдын алу үшін ауруға төзімді бидай сорттары мен линияларын өсіру қажет. Зерттеу жұмысында Түркияның «Күздік бидайды жақсарту бойынша халықаралық орталық»(IWWIP) құрған халықаралық қатты қаракүйе тәлімбағының 75 бидай линиялар жиынтығы

колданылды. Зерттеу жұмысы 2016-2017 жылдары Қазақ егіншілік және өсімдік шаруашылығы ҒЗИ-ның жасанды инфекциялық тәлімбағында жүргізілді. Сезімтал бақылау нұсқасы GEREK 79 генотипі 59,7% масақ қатты қаракүйемен ауырып, сезімталдылықтың жоғары деңгейіне ие болды. Он алты генотип жасанды инфекция аясында ауруға төзімділік танытты. Аурудың ең жоғары деңгейі 59,7%-ға дейін масақтары залалданған бидай линиясында тіркелді. 42 бидай генотипі 2.0-27.3% шамасында қатты қаракүйемен залалданып, орташа төзімді деп танылды. Жасанды жағдайда қатты қаракүйе ауруы бидайдың агрономиялық көрсеткіштеріне әртүрлі әсер етті. Бидай линияларының өнімділік деңгейі 1,13 т/га-дан 7,29 т/га-ға дейін ауытқыды. Болжам жасалынған жоғары оң корреляция коэффициенті дән саны мен дән салмағы ($r = 0,7$), дән салмағы мен 1000 дәннің салмағы ($r = 0,75$) және 1 масақтағы дән салмағы мен жалпы дән салмағы ($r = 0,79$) арасында байқалды. Ал теріс корреляция аурумен залалданған масақтар саны мен басқа агрономиялық көрсеткіштердің арасында байқалды. Идентификацияланған төзімді генотиптер Қазақстандағы қатты қаракүйеге төзімді БИДВАЙ сорттарын құруға арналған селекциялық бағдарламаларда қолдану үшін құнды болып табылады.

Түйін сөздер: бидай, бидай линиялары, қатты қаракүйе, төзімділік, өнімділік, агрономиялық көрсеткіштер.

К. Ж. Тагаев¹, А. И. Моргунов², М. А. Есимбекова³, А. И. Аbugалиева³, Г. О. Баядилова¹

¹Казахский национальный аграрный университет, Алматы, Казахстан,

²Международный центр улучшения кукурузы и пшеницы (СИММИТ), Анкара, Турция,

³Казахский научно-исследовательский институт земледелия и растениеводства,
Алмалыбак, Алматинская область, Казахстан

ВЛИЯНИЕ ТВЕРДОЙ ГОЛОВНИ НА АГРОНОМИЧЕСКИЕ ПОКАЗАТЕЛИ И УСТОЙЧИВОСТЬ ЛИНИИ ОЗИМОЙ ПШЕНИЦЫ

Аннотация. Твердая головня (*Tilletiacaries*) является наиболее опасным грибковым заболеванием пшеницы. Из системы мероприятий по борьбе с твердой головней наиболее радикальной является создание устойчивых сортов и линий пшеницы. В данной работе использованы 75 сортов и линий пшеницы, созданные в «Международном центре по улучшению озимой пшеницы» (IWWIP), Турция. Эксперимент проведен в 2016–2017 гг. в инфекционном стационаре Казахского научно-исследовательского института земледелия и растениеводства. Восприимчивый контроль, GEREK 79, показала восприимчивость (59,7% поврежденных колосьев). Самая высокая уровень болезни составляла 74,4% пораженных колосьев. Шестнадцать генотипов были устойчивыми в условиях искусственного инфицирования. Сорок два генотипов пшеницы были умеренно-устойчивыми, их колосья поражались головней от 2,0% до 27,3%. Агрономические показатели в условиях искусственного инфицирования колебались по-разному по сравнению с контролем. Урожайность генотипов пшеницы колебалась от 1,13 до 7,29 т/га. Ожидаемая высокая положительная корреляция обнаружена между числом зерен и массой зерна ($r = 0,7$), между массой зерна и массой 1000 зерен ($r = 0,75$), а также между массой зерна с главного колоса и общей массой зерна с растения ($r = 0,79$). Отрицательная корреляция обнаружена между колосьями пораженный твердой головней и другими агрономическими признаками. Идентифицированные устойчивые генотипы будут ценными в селекционных программах Казахстана для формирования устойчивых сортов пшеницы к твердой головне.

Ключевые слова: пшеница, линии пшеницы, твердая головня, устойчивость, продуктивность, агрономические признаки.

Information about authors:

Tagayev Kuttymurat Zhurgenbayuly – PhD student of Kazakh national agrarian university, PhD student; kuttymurat.tagayev@mail.ru.

Morgounov Alexey Ivanovich – CIMMIT, the CIMMYT country representative in Turkey; a.morgounov@cgiar.org

Yessimbekova Minura Akhmetovna – Head of Gene pool of Kazakh Research Institute of Agriculture and Plant Growing, Doctor of Biological Sciences; minura.esimbekova@mail.ru.

Abugalieva Aigul Iztelevovna – Head of laboratory biochemistry and grain quality of Kazakh Research Institute of Agriculture and Plant Growing, Doctor of Biological Sciences; kiz_abugalieva@mail.ru.

Bayadilova Gulsun Ongarovana, Associated professor of department Agronomy of Kazakh national agrarian university, Candidate of Biological Sciences; zhalaiirka_kushik@mail.ru.

Publication Ethics and Publication Malpractice in the journals of the National Academy of Sciences of the Republic of Kazakhstan

For information on Ethics in publishing and Ethical guidelines for journal publication see <http://www.elsevier.com/publishingethics> and <http://www.elsevier.com/journal-authors/ethics>.

Submission of an article to the National Academy of Sciences of the Republic of Kazakhstan implies that the described work has not been published previously (except in the form of an abstract or as part of a published lecture or academic thesis or as an electronic preprint, see <http://www.elsevier.com/postingpolicy>), that it is not under consideration for publication elsewhere, that its publication is approved by all authors and tacitly or explicitly by the responsible authorities where the work was carried out, and that, if accepted, it will not be published elsewhere in the same form, in English or in any other language, including electronically without the written consent of the copyright-holder. In particular, translations into English of papers already published in another language are not accepted.

No other forms of scientific misconduct are allowed, such as plagiarism, falsification, fraudulent data, incorrect interpretation of other works, incorrect citations, etc. The National Academy of Sciences of the Republic of Kazakhstan follows the Code of Conduct of the Committee on Publication Ethics (COPE), and follows the COPE Flowcharts for Resolving Cases of Suspected Misconduct (http://publicationethics.org/files/u2/New_Code.pdf). To verify originality, your article may be checked by the Cross Check originality detection service <http://www.elsevier.com/editors/plagdetect>.

The authors are obliged to participate in peer review process and be ready to provide corrections, clarifications, retractions and apologies when needed. All authors of a paper should have significantly contributed to the research.

The reviewers should provide objective judgments and should point out relevant published works which are not yet cited. Reviewed articles should be treated confidentially. The reviewers will be chosen in such a way that there is no conflict of interests with respect to the research, the authors and/or the research funders.

The editors have complete responsibility and authority to reject or accept a paper, and they will only accept a paper when reasonably certain. They will preserve anonymity of reviewers and promote publication of corrections, clarifications, retractions and apologies when needed. The acceptance of a paper automatically implies the copyright transfer to the National Academy of Sciences of the Republic of Kazakhstan.

The Editorial Board of the National Academy of Sciences of the Republic of Kazakhstan will monitor and safeguard publishing ethics.

Правила оформления статьи для публикации в журнале смотреть на сайте:

[www:nauka-nanrk.kz](http://www.nauka-nanrk.kz)

<http://agricultural.kz/>

Редактор *М. С. Ахметова, Т. М. Апендиев, Д. С. Аленов*
Верстка на компьютере *Д. Н. Калкабековой*

Подписано в печать 10.10.2018.
Формат 60x881/8. Бумага офсетная. Печать – ризограф.
5,9 п.л. Тираж 300. Заказ 5.