

ISSN 2224-526X

ҚАЗАҚСТАН РЕСПУБЛИКАСЫ
ҰЛТТЫҚ ҒЫЛЫМ АКАДЕМИЯСЫНЫҢ
Қазақ ұлттық аграрлық университеті

Х А Б А Р Л А Р Ы

ИЗВЕСТИЯ

НАЦИОНАЛЬНОЙ АКАДЕМИИ НАУК
РЕСПУБЛИКИ КАЗАХСТАН
Қазақстан Республикасының
Ғылым Академиясының
Қазақ ұлттық аграрлық университеті

IZVESTIÂ

NATIONAL'NOJ AKADEMII NAUK
RESPUBLIKI KAZAHSTAN
Kazakh national
agrarian university

SERIÂ AGRARNYH NAUK

3 (51)

MAY – JUNE 2019

PUBLISHED SINCE JANUARY 2011

PUBLISHED 6 TIMES A YEAR

ALMATY, NAS RK

Б а с р е д а к т о р

Есполов Т.И.,

э.ғ.д, профессор,

ҚР ҰҒА академигі және вице-президенті

Р е д а к ц и я а л қ а с ы:

Байзақов С.Б., э.ғ.д, проф., ҚР ҰҒА академигі (бас редактордың орынбасары); **Тиреуов К.М.**, э.ғ.д, проф., ҚР ҰҒА академигі (бас редактордың орынбасары); **Елешев Р.Е.**, т.ғ.д, проф., ҚР ҰҒА академигі; **Рау А.Г.**, т.ғ.д, проф., ҚР ҰҒА академигі; **Иванов Н.П.**, в.ғ.д, проф., ҚР ҰҒА академигі; **Кешуов С.А.**, т.ғ.д, проф., ҚР ҰҒА академигі; **Мелдебеков А.**, а.ш.ғ.д., проф., ҚР ҰҒА академигі; **Чоманов У.Ч.**, т.ғ.д, проф., ҚР ҰҒА академигі; **Елюбаев С.З.**, а.ш.ғ.д., проф., ҚР ҰҒА академигі; **Садықұлов Т.**, а.ш.ғ.д., проф., академигі; **Баймұқанов Д.А.**, а.ш.ғ.д., проф., ҚР ҰҒА корр-мүшесі; **Сансызбай А.Р.**, а.ш.ғ.д., проф., ҚР ҰҒА корр-мүшесі; **Умбетаев И.**, а.ш.ғ.д., проф., ҚР ҰҒА академигі; **Оспанов С.Р.**, а.ш.ғ.д., проф., ҚР ҰҒА құрметті мүшесі; **Олейченко С.И.**, а.ш.ғ.д., проф.; **Кененбаев С.Б.**, а.ш.ғ.д., проф., ҚР ҰҒА корр-мүшесі; **Омбаев А.М.**, а.ш.ғ.д., проф. ҚР ҰҒА корр-мүшесі; **Молдашев А.Б.**, э.ғ.д, проф., ҚР ҰҒА құрметті мүшесі; **Сагитов А.О.**, б.ғ.д., ҚР ҰҒА академигі; **Сапаров А.С.**, а.ш.ғ.д., проф., ҚР АШҒА академигі; **Балгабаев Н.Н.**, а.ш.ғ.д., проф.; **Умирзаков С.И.**, т.ғ.д, проф.; **Султанов А.А.**, в.ғ.д, проф., ҚР АШҒА академигі; **Алимкулов Ж.С.**, т.ғ.д., проф., ҚР АШҒА академигі; **Сарсембаева Н.Б.**, в.ғ.д., проф.

Р е д а к ц и я к е н е с і:

Fasler-Kan Elizaveta, Dr., University of Basel Switzerland; **Koolmees Petrus Adrianus**, Prof. Dr., Utrecht University, The Netherlands; **Babadoost-Kondri Mohammad**, Prof., University of Illinois, USA; **Yus Aniza Binti Yusof**, Dr., University Putra, Malaysia; **Hesseln Hayley Fawn**, As. Prof., University of Saskatchewan, Canada; **Alex Morgounov**, Pr., International Maize and Wheat Improvement Center Turkey; **Андреш С.**, Молдова Республикасы ҰҒА академигі; **Гаврилюк Н.Н.**, Украина ҰҒА академигі; **Герасимович Л.С.**, Беларусь Республикасының ҰҒА академигі; **Мамедов Г.**, Азербайжан Республикасының ҰҒА академигі; **Шейко И.П.**, Беларусь Республикасының ҰҒА академигі; **Жалнин Э.В.**, т.ғ.д., проф., Ресей; **Боинчан Б.**, а.ш.ғ.д, проф., Молдова Республикасы; **Юлдашбаев Ю.А.**, а.ш.ғ.д, проф., РФА корр-мүшесі, Ресей.

Главный редактор

Есполов Т.И.,

доктор эконом. наук, проф.,
вице-президент и академик НАН РК

Редакционная коллегия:

Байзаков С.Б., доктор эконом. наук, проф., академик НАН РК (заместитель главного редактора); **Тиреуов К.М.**, доктор эконом. наук, проф., академик НАН РК (заместитель главного редактора); **Елешев Р.Е.**, доктор техн. наук, проф., академик НАН РК; **Рау А.Г.**, доктор техн. наук, проф., академик НАН РК; **Иванов Н.П.**, доктор ветеринар. наук, проф., академик НАН РК; **Кешуов С.А.**, доктор техн. наук, проф., академик НАН РК; **Мелдебеков А.**, доктор сельхоз. наук, проф., академик НАН РК; **Чоманов У.Ч.**, доктор техн. наук, проф., академик НАН РК; **Елюбаев С.З.**, доктор сельхоз. наук, проф., академик НАН РК; **Садыкулов Т.**, доктор сельхоз. наук, проф., академик НАН РК; **Баймуқанов Д.А.**, доктор сельхоз. наук, проф., член-корр. НАН РК; **Сансызбай А.Р.**, доктор сельхоз. наук, проф., член-корр. НАН РК; **Умбетаев И.**, доктор сельхоз. наук, проф., академик НАН РК; **Оспанов С.Р.**, доктор сельхоз. наук, проф., Почетный член НАН РК; **Олейченко С.И.**, доктор сельхоз. наук, проф.; **Кененбаев С.Б.**, доктор сельхоз. наук, проф., член-корр. НАН РК; **Омбаев А.М.**, доктор сельхоз. наук, проф. член-корр. НАН РК.; **Молдашев А.Б.**, доктор эконом. наук, проф., Почетный член НАН РК; **Сагитов А.О.**, доктор биол. наук, академик НАН РК; **Сапаров А.С.**, доктор сельхоз. наук, проф., академик АСХН РК; **Балгабаев Н.Н.**, доктор сельхоз. наук, проф.; **Умирзаков С.И.**, доктор техн. наук, проф.; **Султанов А.А.**, доктор ветеринар. наук, проф., академик АСХН РК; **Алимкулов Ж.С.**, доктор техн. наук, проф., академик АСХН РК; **Сарсембаева Н.Б.**, доктор ветеринар. наук, проф.

Редакционный совет:

Fasler-Kan Elizaveta, Dr., University of asel Switzeland; **Koolmees Petrus Adrianus**, Prof. Dr., Utrecht University, The Netherlands; **Babadoost-Kondri Mohammad**, Prof., University of Illinois, USA; **Yus Aniza Binti Yusof**, Dr., University Putra, Malaysia; **Hesseln Hayley Fawn**, As.Prof., University of Saskatchewan, Canada; **Alex Morgounov**, Pr., International Maize and Wheat Improvement Center Turkey; **Андреш С.**, академик НАН Республики Молдова; **Гаврилюк Н.Н.**, академик НАН Украины; **Герасимович Л.С.**, академик НАН Республики Беларусь; **Мамедов Г.**, академик НАН Республики Азербайджан; **Шейко И.П.**, академик НАН Республики Беларусь; **Жалнин Э.В.**, доктор техн. наук, проф., Россия; **Боинчан Б.**, доктор сельхоз. наук, проф., Республика Молдова; **Юлдашбаев Ю.А.**, доктор сельхоз. наук, проф., член-корр. РАН, Россия.

Известия Национальной академии наук Республики Казахстан. Серия аграрных наук.

ISSN 2224-526X

Собственник: ООО «Национальная академия наук Республики Казахстан» (г. Алматы)

Свидетельство о постановке на учет периодического печатного издания в Комитете информации и архивов Министерства культуры и информации Республики Казахстан № 10895-Ж, выданное 30.04.2010 г.

Периодичность 6 раз в год

Тираж: 300 экземпляров

Адрес редакции: 050010, г. Алматы, ул. Шевченко, 28, ком. 219-220, тел. 272-13-19, 272-13-18

<http://agricultural.kz/index.php/en/>

© Национальная академия наук Республики Казахстан, 2019

Адрес типографии: ИП «Аруна», г. Алматы, ул. Муратбаева, 75

Chief Editor

Espolov T.I.,

Dr. economy. Sciences, prof.,
Vice President and academician of the NAS RK

Editorial Board:

Baizakov S.B., Dr. of economy sciences, prof., academician of NAS RK (deputy editor); **Tireuov K.M.**, Doctor of Economy Sciences., prof., academician of NAS RK (deputy editor); **Eleshev R.E.**, Dr. Of agricultural sciences, prof., academician of NAS RK; **Rau A.G.**, Dr. sciences, prof., academician of NAS RK; **Ivanov N.P.**, Dr. of veterinary sciences, prof., academician of NAS RK; **Keshuov S.A.**, Dr. sciences, prof., academician of NAS RK; **Meldebekov A.**, doctor of agricultural sciences, prof., academician of NAS RK; **Chomanov U.Ch.**, Dr. sciences, prof., academician of NAS RK; **Yelyubayev S.Z.**, Dr. of agricultural sciences, prof., academician of NAS RK; **Sadykulov T.**, Dr. Farm. Sciences, prof., academician of NAS RK; **Baimukanov D.A.**, doctor of agricultural sciences, prof., corresponding member NAS RK; **Sansyzbai A.R.**, doctor of agricultural sciences, prof., corresponding member NAS RK; **Umbetaev I.**, Dr. Farm. Sciences, prof., academician of NAS RK; **Ospanov S.R.**, Dr. agricultural sciences, prof., Honorary Member of NAS RK; **Oleychenko S.N.**, Dr. Of agricultural sciences, prof.; **Kenenbayev S.B.**, Dr. Agricultural sciences, prof., corresponding member NAS RK; **Ombayev A.M.**, Dr. Agricultural sciences, Prof. corresponding member NAS RK; **Moldashev A.B.**, Doctor of Economy sciences, prof., Honorary Member of NAS RK; **Sagitov A.O.**, Dr. biol. sciences, academician of NAS RK; **Saparov A.S.**, Doctor of agricultural sciences, prof., academician of NAS RK; **Balgabaev N.N.**, the doctor agricultural sciences, Prof.; **Umirzakov S.I.**, Dr. Sci. Sciences, Prof.; **Sultanov A.A.**, Dr. of veterinary sciences, prof., academician of the Academy of Agricultural Sciences of Kazakhstan; **Alimkulov J.C.**, Dr. of tekhncial sciences, prof., academician of the Academy of Agricultural sciences of Kazakhstan; **Sarsembayeva N.B.**, Dr. veterinary sciences, prof.

Editorial Board:

Fasler-Kan Elizaveta, Dr., University of Basel Switzzeland; **Koolmees Petrus Adrianus**, Prof. Dr., Utrecht University, The Netherlands; **Babadoost-Kondri Mohammad**, Prof., University of Illinois, USA; **Yus Aniza Binti Yusof**, Dr., University Putra, Malaysia; **Hesseln Hayley Fawn**, As. Prof., University of Saskatchewan, Canada; **Alex Morgounov**, candidate of agricultural sciences, International Maize and Wheat Improvement Center Turkey; **Andresh S.**, academician of NAS of Moldova; **Gavriluk N.N.**, academician of NAS of Ukraine; **Gerasimovich L.S.**, academician of NAS of Belorassia; **Mamadov G.**, academician of NAS of Azerbaijan; **Sheiko I.P.**, academician of NAS of Belorassia; **Zhalnin E.V.**, Dr. of technical sciences, professor, Russia, **Boinchan B.**, doctor of agricultural sciences, prof., Moldova; **Yuldashbayev Y.A.**, doctor of agricultural sciences, prof., corresponding member of RAS, Russia.

News of the National Academy of Sciences of the Republic of Kazakhstan. Series of Agrarian Sciences.

ISSN 2224-526X

Owner: RPA "National Academy of Sciences of the Republic of Kazakhstan" (Almaty)

The certificate of registration of a periodic printed publication in the Committee of Information and Archives of the Ministry of Culture and Information of the Republic of Kazakhstan N 10895-Ж, issued 30.04.2010

Periodicity: 6 times a year

Circulation: 300 copies

Editorial address: 28, Shevchenko str., of.219-220, Almaty, 050010, tel. 272-13-19, 272-13-18,
<http://nauka-nanrk.kz/agricultural.kz>

© National Academy of Sciences of the Republic of Kazakhstan, 2019

Address of printing house: ST "Aruna", 75, Muratbayev str, Almaty

NEWS

OF THE NATIONAL ACADEMY OF SCIENCES OF THE REPUBLIC OF KAZAKHSTAN

SERIES OF AGRICULTURAL SCIENCES

ISSN 2224-526X

Volume 3, Number 51 (2019), 11 – 15

<https://doi.org/10.32014/2019.2224-526X.28>

UDK 619:616.3

T. K. Boranbayeva, Zh. K. Tulemissova, D. M. Khussainov

Kazakh National Agrarian University, Almaty, Kazakhstan.

E-mail: bor-tog@mail.ru, zhanara.tulemissova@gmail.com, doctor-vet@mail.ru

**STUDYING THE ANTAGONISTIC PROPERTIES
OF LACTOBACTERIN-TK² PROBIOTICS
ON PATHOGENIC STRAINS CAUSING GASTROINTESTINAL
PATHOLOGY IN CALVES AND LAMPS**

Abstract. This article contains the researches of antagonistic activity of pathogenic bacteria causing a gastrointestinal disorder in calves and lambs in their first days of life. The antagonistic activity on the pathogenic intestinal microflora has been studied *in vitro*.

Key words: *Escherichia Coli*, *Salmonella AbortusOvis*, *Proteus Vulgaris*, Lactobacterin-TK², calves and lambs, gastrointestinal pathology, research.

Introduction. Recently, among infectious diseases of calves and lambs, gastrointestinal diseases of bacterial etiology occupy a special place.

Pathogenic strains of *Escherichia*, *Salmonella* and *Proteus*, which cause gastrointestinal disorder in calves and lambs in their first days of life, play a leading role in the nosological profile of these diseases [1].

Pathogenic strains of *Escherichia Coli*, *Salmonella AbortusOvis* and *Proteus Vulgaris* belong to Enterobacteriaceae family and, by morphology, are small gram-negative straight rods [2].

According to the official definition, escherichiosis or colibacteriosis in animals is an acutely occurring zoonotic disease, which usually infests young animals, and characterized by diarrhea, septicemia, toxemia and enteritis, general dehydration of the body, disorders of the central nervous system, symptoms of gradually increasing depression and weakness.

Escherichia are permanent inhabitants of the intestines of warm-blooded animals, generally, they live in the lower parts of the gastrointestinal tract. Some of them can cause lesions of the gastrointestinal tract that was experimentally proven by G. N. Gabrichevsky in 1894 and confirmed clinically by A. Adam. Calves and lambs become ill with escherichiosis mainly in the first 2-7 days of life, but also the disease often affects one- and two-month-old calves and lambs [3]. The incidence of disease can reach up to 90%, and the mortality rate is about 30-50%. The incubation period of the disease lasts from several hours to 1-2 days. Escherichiosis in calves and lambs can occur in the septic, enteritic and enterotoxemic forms [4].

A characteristic symptom of the disease is the feces color, which in most cases has a white, yellowish or bright yellowish color with a greenish tinge and a sour odor. In calves and lambs affected by escherichiosis, subfebrile temperature reaches 40-40.5°C, the rapid toxicosis development is manifested by the rejection of colostrum, lethargy and lying long. [5]

Salmonellosis is an infectious disease in young farm animals, which is characterized, when in acute form, by fever, septicemia and diarrhea, and when in chronic form – by pneumonia.

The sources of pathogen of this infection are sick and ill animals. Adult animals can be salmonella carriers, releasing the pathogen with milk and feces, abortus fetus, amniotic fluid, and outflows from the birth canal. Sick young animals produce a pathogen with feces, urine, nasal effusion, saliva [6].

In case of delayed medicated treatment, the condition of sick animals worsens by the end of the first or second day of disease and is characterized by a complete lack of appetite and sucking reflex, adynamia, anuria, a decrease in body temperature to 36-37.7°C, cyanosis and dry mucous membranes, deep drooping of the eyeballs, involuntary outflow of watery feces from the anus. Calves and lambs predominantly die from dehydration.

Proteas are one of the most well-known genera of the Enterobacteriaceae family.

All types of proteas belong to the group of conditionally pathogenic microorganisms, which exhibit a negative effect on the body while reducing antimicrobial protection. For the development of infection, the virulent properties of bacteria are important. The most important pathogenic factors of the proteas are: fimbriae, bacterial proteas and urease, hemolysins, hemagglutinins, and the ability to “swarm”.

Dysbiotic disorders of the gastrointestinal tract can cause acute intestinal diseases in newborn calves and lambs due to the prevalence of enterobacteria over the symbiotic flora, that is a feature in the development of intestinal microbiocenosis in calves and lambs in the first 7 days of life [7].

Enteropathogenic bacteria are able to show resistance to certain types of antibiotics. Recently, one of the important advances in the field of medicine is the establishment of the fact that the microflora inhabiting the organism of animals is not beneficial only, but necessary for its vital activity as well. It is known that the most numerous and complex in terms of its composition is the bacteria population in the intestine, especially in its lower parts [8].

Therefore, the search for effective forms of prophylaxis, using the beneficial microflora, which are able to stop the spread and development of dysbiotic conditions of the gastrointestinal tract of calves and lambs in time, is a hot topic.

Relevance of research. In our country, gastrointestinal diseases of bacterial etiology in calves and lambs remain relevant and continue to cause significant economic damage, since without introducing new means of prevention and therapy of these diseases it will be impossible to obtain high-quality animal products [2].

These pathologies also affect the normal gastrointestinal microflora of the animal organism. A number of scientists believe that one of the stimulating factors for occurrence of gastrointestinal pathologies contributes to the disorders in the gastrointestinal microbiocenosis. These disorders causing this series of disorders are united by a common name - dysbacteriosis [9].

A number of authors believe that the cause of dysbacteriosis is the excessive use of drugs, in particular antibiotics and other antimicrobial drugs. Their accumulation in the body, especially in young animals, leads to a weakening of natural resistance and, as a rule, leads to the formation of infectious processes [1, 3, 9, 10].

It is well known that probiotics are one of the most promising agents for treatment and prevention of dysbiotic conditions. Probiotic drugs or food products contain live microorganisms, most often lactobacteria (lactobacilli, bifidobacteria, enterococci). World experience shows that in the prevention and treatment of gastrointestinal diseases in young animals, replacement therapy is important, which is aimed at restoring the intestinal biocenosis through the regular introduction of live bacteria, representatives of the normal intestinal microflora. The drugs they contain are known as probiotics. Probiotics are used to stimulate the immunity, prevent and treat mixed gastrointestinal infections, digestive disorders resulting from a sudden change in the diet composition, disturbed feeding regimes, technological stress, etc. [1, 11].

Currently, many farms successfully practice of calf growing [12]. The advantage of probiotic agents containing lactobacilli is that they are harmless to the organism and there is no addiction to them during prolonged use, there are no side effects completely [13].

Purpose and methods of research. Antagonistic activity of associations of probiotic bacteria was determined by co-cultivation in relation to the culture of *Escherichia Coli*, *Salmonella Abortus Ovis* and *Proteus Vulgaris*.

On dense nutrient media, these pathogens form round colonies of 2-4 mm in diameter, with a smooth, convex surface and a smooth edge. *Escherichia* on MPA (Meat-Peptide Agar) - the colonies are translucent, grayish. On Levin's medium – they are dark blue, purple, black with or without metallic luster; on Mc-Conkey medium - pink, red (some strains of *Escherichia* may not ferment lactose and form colorless colonies on the listed media). Proteas and colonies of pathological microorganisms of other serogroups on BGA (Brilliant Green Agar) form pale pink or red-crimson colonies, transparent, surrounded by a brilliant



a) *Salmonella Abortus Ovis*
and *Proteus Vulgaris*

b) Stains of *Escherichia*, *Salmonella*
Abortus Ovis and *Proteus Vulgaris*

c) Stain
of *Escherichia Coli*


Figure 1 – Colonies of *Escherichia Coli*, *Salmonella Abortus Ovis* and *Proteus Vulgaris* on a dense nutrient medium

red halo. *Salmonella* on BSA (Bismuth Sulphite Agar) usually form black colonies with metallic luster, surrounded by blackening as a result of hydrogen sulfide production and recovery of sulphite to ferrous sulphide, which is black.

For the researches, the deposited passported lactobacillus strain, *Escherichia Coli*, *Salmonella Abortus Ovis* and *Proteus Vulgaris*, contained in Lactobacterin-TK², were used. Cultivation of the *Lactobacillus acidophilus* B-RKM-01511 strain of probiotic Lactobacterin-TK² was carried out under aerobic conditions at a temperature of 37°C for 24-48 hours in a milk hydrolyzate medium. Enterobacteria were cultured under aerobic conditions at 37°C for 24 hours in Müller Hinton agar medium.

A diffusion well method was used. Separately, 1 ml of cultures of *Escherichia Coli*, *Salmonella Abortus Ovis* and *Proteus Vulgaris* grown within 24 hours were added to sterile Petri dishes, having a titer of 10⁵ microbial cells per ml according to the turbidity standard for opportunistic strains of these bacteria, and then 20 ml of molten and cooled to 40-45°C MPA. When the coating of dishes has hardened in the form of metal stamp, wells with a diameter of 10 mm were cut out, and 100 µl of probiotic bacteria associations, which are part of Lactobacterin-TK², were introduced thereinto. After incubation at room temperature, the dishes were placed in a thermostat (37°C) for 24-48 hours. Then, the diameter of growth inhibition areas of tested microorganisms around the well, including its diameter, was determined.

The results of research of the probiotic activity of Lactobacterin-TK² are shown in table.



a) *Escherichia Coli*,

b) *Salmonella Abortus Ovis*

c) *Proteus Vulgaris*,

Figure 2 – Determination by the diffusion method of the well diameters of growth inhibition areas of tested microorganisms

Zone of growth inhibition of the antagonistic activity of the culture of probiotic bacteria
Lactobacillus acidophilus B-RKM-01511 against *Escherichia Coli*, *Salmonella Abortus Ovis* and *Proteus Vulgaris*

Culture	Antagonistic activity of the probiotic drug (growth inhibition zone in mm)		
	Lactobacillus acidophilus B-RKM-01511		
	Sensibility	Dose-dependent zone	Resistance
<i>Escherichia Coli</i>	≥ 22	20-21	≤ 19
<i>Salmonella Abortus Ovis</i>	≥ 26	23,5-25	$\leq 22,5$
<i>Proteus Vulgaris</i>	≥ 21	20-20	≤ 19

From the data in table it can be seen that the probiotic strain of Lactobacterin-TK²- *Lactobacillus acidophilus* B-RKM-01511 has antagonistic activity to strains *Escherichia Coli*, *Salmonella Abortus Ovis* and *Proteus Vulgaris*.

It was found that the strains, which are part of probiotic Lactobacterin-TK² are able to show antagonism with regard to *Escherichia Coli*, *Salmonella Abortus Ovis* and *Proteus Vulgaris* that determines the high efficacy of the drug.

Results. Made researches have shown that Lactobacterin-TK² probiotic strain – *Lactobacillus acidophilus* B-RKM-01511 has a high antagonistic activity and is able to succeed in the gastrointestinal tract of calves and lambs. The results can serve as a basis for the inclusion of Lactobacterin-TK² in the scheme for complex use in the prevention of dysbiotic conditions of the gastrointestinal tract in sick animals.

Т. К. Боранбаева, Ж. К. Тулемисова, Д. М. Хусаинов

Қазақ ұлттық аграрлық университеті, Алматы, Қазақстан

**БҰЗАУЛАР МЕН ҚОЗЫЛАРДЫҢ АСҚАЗАН-ІШЕК ЖОЛЫ
ПАТОЛОГИЯСЫН ТУДЫРАТЫН ПАТОГЕНДІ ШТАМДАРҒА ҚАРСЫ
"ЛАКТОБАКТЕРИН-ТК²" ПРОБИОТИГИНІҢ
АНТАГОНИСТІК ҚАСИЕТТЕРІН ЗЕРТТЕУ**

Аннотация. Мақалада жана туылған бұзаулар мен қозылардың асқазан-ішек жолы қызметінің бұзылуына әкелетін патогенді бактериялардың антагонистік белсенділігі зерттелген. In vitro жағдайында ішектің патогенді микрофлорасына қарсы антагонистік белсенділік зерттелді.

Түйін сөздер: *Escherichia Coli*, *Salmonella Abortus Ovis*, *Proteus Vulgaris*, Lactobacterin-TK², бұзау және қозылар, асқазан-ішек патологиясы, зерттеу.

Т. К. Боранбаева, Ж. К. Тулемисова, Д. М. Хусаинов

Казахский национальный аграрный университет, Алматы, Казахстан

**ИЗУЧЕНИЕ АНТАГОНИСТИЧЕСКИХ СВОЙСТВ
ПРОБИОТИКА «ЛАКТОБАКТЕРИН-ТК²» НА ПАТОГЕННЫЕ ШТАММЫ
ВЫЗЫВАЮЩИЕ ЖЕЛУДОЧНО-КИШЕЧНУЮ ПАТОЛОГИЮ
У ТЕЛЯТ И ЯГНЯТ**

Аннотация. В статье изучена антагонистическая активность патогенных бактерий, вызывающих расстройство деятельности желудочно-кишечного тракта у телят и ягнят в первые дни жизни. В условиях *in vitro* изучена антагонистическая активность на патогенную микрофлору кишечника.

Ключевые слова: *Escherichia Coli*, *Salmonella Abortus Ovis*, *Proteus Vulgaris*, препарат «Лактобактерин-ТК²», телята и ягнята, желудочно-кишечная патология, исследования.

Information about authors:

Boranbayeva Togzhan, Kazakh National Agrarian University, Almaty, Kazakhstan; bor-tog@mail.ru; <https://orcid.org/0000-0002-1159-1200>

Tulemissova Zhanara, Kazakh National Agrarian University, Almaty, Kazakhstan; zhanara.tulemissova@gmail.com; <https://orcid.org/0000-0002-6894-7386>

Khussainov Damir, Kazakh National Agrarian University, Almaty, Kazakhstan; doctor-vet@mail.ru; <https://orcid.org/0000-0002-0244-8381>

REFERENCES

[1] Suvorov A., Gromova L., Borshev Y., Ermolenko E., Karaseva A., Gruzdkov A. Influence of probiotic enterococci on the gastrointestinal tract of rats before and after the treatment of antibiotic associated dysbiosis // The Joint Meeting of The XVIIth International Symposium on Gnotobiology and The XXXIVth Congress of the Society for Microbial Ecology and Disease Navios Yokohama. Japan: Yokohama, 2011. P. 108.

[2] Bondarenko V.M., Rubakova Je.I., Lavrova V.A. Immunostimulirujushhee dejstvie laktobakterij, ispol'zuemyh v kachestve osnovy preparatov probiotikov // Mikrobiol. zhurn. 1998. N 4. P. 107-111.

[3] Gaggia F., Mattarelli P., Biavati B. Probiotics and prebiotics in animal feeding for safe food production // International Journal of Food Microbiology. 2010. Vol. 141. P. 15-28.

[4] Gavrilova N.N., Ratnikova I.A., Bajakysheva K., Utegenova N.M., Belikova O.A. Sposobnost' molochnokislyh bakterij probiotika polilaktovit sintezirovat' biologicheski aktivnye veshhestva // Izvestija NAN RK. Serija biologicheskaja i medicinskaja. 2017. N 6. P. 112-119. ISSN 2518-1629 (Online), ISSN 2224-5308 (Print). <https://doi.org/10.32014/2018.25181629>

[5] Tulemissova Zhet al. Prophylaxy of gasstro-intestinal diseases of young animals // Jour. of Anim. and Vet. Ad. 2013. Vol. 12. P. 1645-1650.

[6] Zhang X.S., Chase-Topping M.E., McKendrick I.J., Savill N.J., Woolhouse M.E.J. Spread of E. coli O157 infection among Scottish cattle farms: stochastic models and model selection // Epidemics. 2010. Vol. 2, N 1. P. 11-20.

[7] Vlková E., Rada V., Trojanová I., Killer J., Smehilová M., Molatová Z. Occurrence of bifidobacteria in faeces of calves fed milk or a combined diet // Arch Anim Nutr. 2008. Vol. 62, N 5. P. 359-365.

[8] García-Hernández Y. et al. Isolation, characterization and evaluation of probiotic lactic acid bacteria for potential use in animal production // Res. Vet. Sci. 2016. Vol. 108. P. 125-132.

[9] Von Baum H., Marre R. Antimicrobial resistance of Escherichia coli and therapeutic implications // Int. J. Med. Microbiol. 2005. Vol. 295. P. 503-511.

[10] Ripamonti B. et al. Screening of species-specific lactic acid bacteria for veal calves multi-strain probiotic adjuncts // Anaerobe. 2011. Vol. 17, N 3. P. 97-105.

[11] Bayatkouhsar J., Tahmasebi A.M., Naserian A.A., Mokarram R.R., Valizadeh R. Effects of supplementation of lactic acid bacteria on growth performance, blood metabolites and fecal coliform and lactobacilli of young dairy calves // Anim. Feed Sci. Technol. 2013. Vol. 186, N 1-2. P. 1-11.

[12] Semenov V.G. et al. Activation of adaptogenesis and bioresource potential of calves under the conditions of traditional and adaptive technologies // Bulletin of NAS of RK. 2019. Vol. 1, N 377. P. 175-189. 2518-1467 (Online), ISSN 1991-3494 (Print). <https://doi.org/10.32014/2019.2518-1467.20>

[13] Masucci F., De Rosa G., Grasso F., Napolitano F., Esposito G., Di Francia A. Performance and immune response of buffalo calves supplemented with probiotic // Livest. Sci. 2011. Vol. 137, N 1-3. P. 24-30.

Publication Ethics and Publication Malpractice in the journals of the National Academy of Sciences of the Republic of Kazakhstan

For information on Ethics in publishing and Ethical guidelines for journal publication see <http://www.elsevier.com/publishingethics> and <http://www.elsevier.com/journal-authors/ethics>.

Submission of an article to the National Academy of Sciences of the Republic of Kazakhstan implies that the described work has not been published previously (except in the form of an abstract or as part of a published lecture or academic thesis or as an electronic preprint, see <http://www.elsevier.com/postingpolicy>), that it is not under consideration for publication elsewhere, that its publication is approved by all authors and tacitly or explicitly by the responsible authorities where the work was carried out, and that, if accepted, it will not be published elsewhere in the same form, in English or in any other language, including electronically without the written consent of the copyright-holder. In particular, translations into English of papers already published in another language are not accepted.

No other forms of scientific misconduct are allowed, such as plagiarism, falsification, fraudulent data, incorrect interpretation of other works, incorrect citations, etc. The National Academy of Sciences of the Republic of Kazakhstan follows the Code of Conduct of the Committee on Publication Ethics (COPE), and follows the COPE Flowcharts for Resolving Cases of Suspected Misconduct (http://publicationethics.org/files/u2/New_Code.pdf). To verify originality, your article may be checked by the Cross Check originality detection service <http://www.elsevier.com/editors/plagdetect>.

The authors are obliged to participate in peer review process and be ready to provide corrections, clarifications, retractions and apologies when needed. All authors of a paper should have significantly contributed to the research.

The reviewers should provide objective judgments and should point out relevant published works which are not yet cited. Reviewed articles should be treated confidentially. The reviewers will be chosen in such a way that there is no conflict of interests with respect to the research, the authors and/or the research funders.

The editors have complete responsibility and authority to reject or accept a paper, and they will only accept a paper when reasonably certain. They will preserve anonymity of reviewers and promote publication of corrections, clarifications, retractions and apologies when needed. The acceptance of a paper automatically implies the copyright transfer to the National Academy of Sciences of the Republic of Kazakhstan.

The Editorial Board of the National Academy of Sciences of the Republic of Kazakhstan will monitor and safeguard publishing ethics.

Правила оформления статьи для публикации в журнале смотреть на сайте:

www:nauka-nanrk.kz

<http://agricultural.kz/index.php/en/>

Редактор *М. С. Ахметова, Т. М. Апендиев, Д. С. Аленов*
Верстка на компьютере *Д. Н. Калкабековой*

Подписано в печать 11.06.2019.
Формат 60x881/8. Бумага офсетная. Печать – ризограф.
5,75 п.л. Тираж 300. Заказ 3.