

ISSN 2224-526X

ҚАЗАҚСТАН РЕСПУБЛИКАСЫ
ҰЛТТЫҚ ҒЫЛЫМ АКАДЕМИЯСЫНЫҢ
Қазақ ұлттық аграрлық университеті

Х А Б А Р Л А Р Ы

ИЗВЕСТИЯ

НАЦИОНАЛЬНОЙ АКАДЕМИИ НАУК
РЕСПУБЛИКИ КАЗАХСТАН
Қазақстан Республикасының
Ғылым Академиясының
Қазақ ұлттық аграрлық университеті

IZVESTIÂ

NATIONAL'NOJ AKADEMII NAUK
RESPUBLIKI KAZAHSTAN
Kazakh national
agrarian university

SERIÂ AGRARNYH NAUK

3 (51)

MAY – JUNE 2019

PUBLISHED SINCE JANUARY 2011

PUBLISHED 6 TIMES A YEAR

ALMATY, NAS RK

Б а с р е д а к т о р

Есполов Т.И.,

э.ғ.д, профессор,

ҚР ҰҒА академигі және вице-президенті

Р е д а к ц и я а л қ а с ы:

Байзақов С.Б., э.ғ.д, проф., ҚР ҰҒА академигі (бас редактордың орынбасары); **Тиреуов К.М.**, э.ғ.д, проф., ҚР ҰҒА академигі (бас редактордың орынбасары); **Елешев Р.Е.**, т.ғ.д, проф., ҚР ҰҒА академигі; **Рау А.Г.**, т.ғ.д, проф., ҚР ҰҒА академигі; **Иванов Н.П.**, в.ғ.д, проф., ҚР ҰҒА академигі; **Кешуов С.А.**, т.ғ.д, проф., ҚР ҰҒА академигі; **Мелдебеков А.**, а.ш.ғ.д, проф., ҚР ҰҒА академигі; **Чоманов У.Ч.**, т.ғ.д, проф., ҚР ҰҒА академигі; **Елюбаев С.З.**, а.ш.ғ.д, проф., ҚР ҰҒА академигі; **Садықұлов Т.**, а.ш.ғ.д, проф., академигі; **Баймұқанов Д.А.**, а.ш.ғ.д, проф., ҚР ҰҒА корр-мүшесі; **Сансызбай А.Р.**, а.ш.ғ.д, проф., ҚР ҰҒА корр-мүшесі; **Умбетаев И.**, а.ш.ғ.д, проф., ҚР ҰҒА академигі; **Оспанов С.Р.**, а.ш.ғ.д, проф., ҚР ҰҒА құрметті мүшесі; **Олейченко С.И.**, а.ш.ғ.д, проф.; **Кененбаев С.Б.**, а.ш.ғ.д, проф., ҚР ҰҒА корр-мүшесі; **Омбаев А.М.**, а.ш.ғ.д, проф. ҚР ҰҒА корр-мүшесі; **Молдашев А.Б.**, э.ғ.д, проф., ҚР ҰҒА құрметті мүшесі; **Сагитов А.О.**, б.ғ.д, ҚР ҰҒА академигі; **Сапаров А.С.**, а.ш.ғ.д, проф., ҚР АШҒА академигі; **Балгабаев Н.Н.**, а.ш.ғ.д, проф.; **Умирзаков С.И.**, т.ғ.д, проф.; **Султанов А.А.**, в.ғ.д, проф., ҚР АШҒА академигі; **Алимкулов Ж.С.**, т.ғ.д, проф., ҚР АШҒА академигі; **Сарсембаева Н.Б.**, в.ғ.д, проф.

Р е д а к ц и я к е н е с і:

Fasler-Kan Elizaveta, Dr., University of Basel Switzerland; **Koolmees Petrus Adrianus**, Prof. Dr., Utrecht University, The Netherlands; **Babadoost-Kondri Mohammad**, Prof., University of Illinois, USA; **Yus Aniza Binti Yusof**, Dr., University Putra, Malaysia; **Hesseln Hayley Fawn**, As. Prof., University of Saskatchewan, Canada; **Alex Morgounov**, Pr., International Maize and Wheat Improvement Center Turkey; **Андреш С.**, Молдова Республикасы ҰҒА академигі; **Гаврилюк Н.Н.**, Украина ҰҒА академигі; **Герасимович Л.С.**, Беларусь Республикасының ҰҒА академигі; **Мамедов Г.**, Азербайжан Республикасының ҰҒА академигі; **Шейко И.П.**, Беларусь Республикасының ҰҒА академигі; **Жалнин Э.В.**, т.ғ.д, проф., Ресей; **Боинчан Б.**, а.ш.ғ.д, проф., Молдова Республикасы; **Юлдашбаев Ю.А.**, а.ш.ғ.д, проф., РФА корр-мүшесі, Ресей.

Главный редактор

Есполов Т.И.,

доктор эконом. наук, проф.,
вице-президент и академик НАН РК

Редакционная коллегия:

Байзаков С.Б., доктор эконом. наук, проф., академик НАН РК (заместитель главного редактора); **Тиреуов К.М.**, доктор эконом. наук, проф., академик НАН РК (заместитель главного редактора); **Елешев Р.Е.**, доктор техн. наук, проф., академик НАН РК; **Рау А.Г.**, доктор техн. наук, проф., академик НАН РК; **Иванов Н.П.**, доктор ветеринар. наук, проф., академик НАН РК; **Кешуов С.А.**, доктор техн. наук, проф., академик НАН РК; **Мелдебеков А.**, доктор сельхоз. наук, проф., академик НАН РК; **Чоманов У.Ч.**, доктор техн. наук, проф., академик НАН РК; **Елюбаев С.З.**, доктор сельхоз. наук, проф., академик НАН РК; **Садыкулов Т.**, доктор сельхоз. наук, проф., академик НАН РК; **Баймуқанов Д.А.**, доктор сельхоз. наук, проф., член-корр. НАН РК; **Сансызбай А.Р.**, доктор сельхоз. наук, проф., член-корр. НАН РК; **Умбетаев И.**, доктор сельхоз. наук, проф., академик НАН РК; **Оспанов С.Р.**, доктор сельхоз. наук, проф., Почетный член НАН РК; **Олейченко С.И.**, доктор сельхоз. наук, проф.; **Кененбаев С.Б.**, доктор сельхоз. наук, проф., член-корр. НАН РК; **Омбаев А.М.**, доктор сельхоз. наук, проф. член-корр. НАН РК.; **Молдашев А.Б.**, доктор эконом. наук, проф., Почетный член НАН РК; **Сагитов А.О.**, доктор биол. наук, академик НАН РК; **Сапаров А.С.**, доктор сельхоз. наук, проф., академик АСХН РК; **Балгабаев Н.Н.**, доктор сельхоз. наук, проф.; **Умирзаков С.И.**, доктор техн. наук, проф.; **Султанов А.А.**, доктор ветеринар. наук, проф., академик АСХН РК; **Алимкулов Ж.С.**, доктор техн. наук, проф., академик АСХН РК; **Сарсембаева Н.Б.**, доктор ветеринар. наук, проф.

Редакционный совет:

Fasler-Kan Elizaveta, Dr., University of asel Switzeland; **Koolmees Petrus Adrianus**, Prof. Dr., Utrecht University, The Netherlands; **Babadoost-Kondri Mohammad**, Prof., University of Illinois, USA; **Yus Aniza Binti Yusof**, Dr., University Putra, Malaysia; **Hesseln Hayley Fawn**, As.Prof., University of Saskatchewan, Canada; **Alex Morgounov**, Pr., International Maize and Wheat Improvement Center Turkey; **Андреш С.**, академик НАН Республики Молдова; **Гаврилюк Н.Н.**, академик НАН Украины; **Герасимович Л.С.**, академик НАН Республики Беларусь; **Мамедов Г.**, академик НАН Республики Азербайджан; **Шейко И.П.**, академик НАН Республики Беларусь; **Жалнин Э.В.**, доктор техн. наук, проф., Россия; **Боинчан Б.**, доктор сельхоз. наук, проф., Республика Молдова; **Юлдашбаев Ю.А.**, доктор сельхоз. наук, проф., член-корр. РАН, Россия.

Известия Национальной академии наук Республики Казахстан. Серия аграрных наук.

ISSN 2224-526X

Собственник: ООО «Национальная академия наук Республики Казахстан» (г. Алматы)

Свидетельство о постановке на учет периодического печатного издания в Комитете информации и архивов Министерства культуры и информации Республики Казахстан № 10895-Ж, выданное 30.04.2010 г.

Периодичность 6 раз в год

Тираж: 300 экземпляров

Адрес редакции: 050010, г. Алматы, ул. Шевченко, 28, ком. 219-220, тел. 272-13-19, 272-13-18

<http://agricultural.kz/index.php/en/>

© Национальная академия наук Республики Казахстан, 2019

Адрес типографии: ИП «Аруна», г. Алматы, ул. Муратбаева, 75

Chief Editor

Espolov T.I.,

Dr. economy. Sciences, prof.,
Vice President and academician of the NAS RK

Editorial Board:

Baizakov S.B., Dr. of economy sciences, prof., academician of NAS RK (deputy editor); **Tireuov K.M.**, Doctor of Economy Sciences., prof., academician of NAS RK (deputy editor); **Eleshev R.E.**, Dr. Of agricultural sciences, prof., academician of NAS RK; **Rau A.G.**, Dr. sciences, prof., academician of NAS RK; **Ivanov N.P.**, Dr. of veterinary sciences, prof., academician of NAS RK; **Keshuov S.A.**, Dr. sciences, prof., academician of NAS RK; **Meldebekov A.**, doctor of agricultural sciences, prof., academician of NAS RK; **Chomanov U.Ch.**, Dr. sciences, prof., academician of NAS RK; **Yelyubayev S.Z.**, Dr. of agricultural sciences, prof., academician of NAS RK; **Sadykulov T.**, Dr. Farm. Sciences, prof., academician of NAS RK; **Baimukanov D.A.**, doctor of agricultural sciences, prof., corresponding member NAS RK; **Sansyzbai A.R.**, doctor of agricultural sciences, prof., corresponding member NAS RK; **Umbetaev I.**, Dr. Farm. Sciences, prof., academician of NAS RK; **Ospanov S.R.**, Dr. agricultural sciences, prof., Honorary Member of NAS RK; **Oleychenko S.N.**, Dr. Of agricultural sciences, prof.; **Kenenbayev S.B.**, Dr. Agricultural sciences, prof., corresponding member NAS RK; **Ombayev A.M.**, Dr. Agricultural sciences, Prof. corresponding member NAS RK; **Moldashev A.B.**, Doctor of Economy sciences, prof., Honorary Member of NAS RK; **Sagitov A.O.**, Dr. biol. sciences, academician of NAS RK; **Saparov A.S.**, Doctor of agricultural sciences, prof., academician of NAS RK; **Balgabaev N.N.**, the doctor agricultural sciences, Prof.; **Umirzakov S.I.**, Dr. Sci. Sciences, Prof.; **Sultanov A.A.**, Dr. of veterinary sciences, prof., academician of the Academy of Agricultural Sciences of Kazakhstan; **Alimkulov J.C.**, Dr. of tekhncial sciences, prof., academician of the Academy of Agricultural sciences of Kazakhstan; **Sarsembayeva N.B.**, Dr. veterinary sciences, prof.

Editorial Board:

Fasler-Kan Elizaveta, Dr., University of Basel Switzzeland; **Koolmees Petrus Adrianus**, Prof. Dr., Utrecht University, The Netherlands; **Babadoost-Kondri Mohammad**, Prof., University of Illinois, USA; **Yus Aniza Binti Yusof**, Dr., University Putra, Malaysia; **Hesseln Hayley Fawn**, As. Prof., University of Saskatchewan, Canada; **Alex Morgounov**, candidate of agricultural sciences, International Maize and Wheat Improvement Center Turkey; **Andresh S.**, academician of NAS of Moldova; **Gavriluk N.N.**, academician of NAS of Ukraine; **Gerasimovich L.S.**, academician of NAS of Belorassia; **Mamadov G.**, academician of NAS of Azerbaijan; **Sheiko I.P.**, academician of NAS of Belorassia; **Zhalnin E.V.**, Dr. of technical sciences, professor, Russia, **Boinchan B.**, doctor of agricultural sciences, prof., Moldova; **Yuldashbayev Y.A.**, doctor of agricultural sciences, prof., corresponding member of RAS, Russia.

News of the National Academy of Sciences of the Republic of Kazakhstan. Series of Agrarian Sciences.

ISSN 2224-526X

Owner: RPA "National Academy of Sciences of the Republic of Kazakhstan" (Almaty)

The certificate of registration of a periodic printed publication in the Committee of Information and Archives of the Ministry of Culture and Information of the Republic of Kazakhstan N 10895-Ж, issued 30.04.2010

Periodicity: 6 times a year

Circulation: 300 copies

Editorial address: 28, Shevchenko str., of.219-220, Almaty, 050010, tel. 272-13-19, 272-13-18,
<http://nauka-nanrk.kz/agricultural.kz>

© National Academy of Sciences of the Republic of Kazakhstan, 2019

Address of printing house: ST "Aruna", 75, Muratbayev str, Almaty

NEWS

OF THE NATIONAL ACADEMY OF SCIENCES OF THE REPUBLIC OF KAZAKHSTAN

SERIES OF AGRICULTURAL SCIENCES

ISSN 2224-526X

Volume 3, Number 51 (2019), 51 – 58

<https://doi.org/10.32014/2019.2224-526X.35>

UDC 635,658: 632.11

A. Zh. Saykenova¹, T. N. Nurgasenov¹,
M. S. Kudaibergenov², S. V. Didorenko², B. R. Saykenov¹

¹Kazakh National Agrarian University, Almaty, Kazakhstan,

²Kazakh Research Institute of Agriculture and Crop Production, Almaty region, Kazakhstan

COMPARATIVE EVALUATION OF DROUGHTABILITY OF VARIETIES OF LENTILES

Abstract. Article results in laboratory experiments to determine the relative drought variety samples method lentil seed germination in the sand in a solution of PEG -10%, 15%. Data on drought tolerance are given. Analysis of lentil samples under conditions of osmotic stress (PEG 10% and PEG 15%) revealed highly dry-resistant samples: K-6, 39113 (Russia).

Keywords: lentils, variety samples, germination, drought resistance, osmotic stress.

Introduction. Lentils are among the most valuable food leguminous crops grown mainly for grain, which is more than a third of protein. Lentil protein contains essential amino acids for the body. Lentil dishes serve for us as suppliers of essential vitamins and minerals that are completely absorbed by the human body. On the content of iron, for example, it has no equal. Lentil has another very valuable property; it does not accumulate in itself any harmful or toxic elements (nitrates, radionuclide's, etc.). Due to this, lentils grown anywhere in the world can be considered an environmentally friendly product [1].

In Kazakhstan, lentils are sown on the territory of the order of 331,566.5 thousand hectares, mostly these plots are located in the north of the republic and are grown in non-irrigated conditions.

In recent years the numbers of dry years are increasing due to the sharp warming on the planet. Drought is a rather frequent phenomenon that disrupts the normal course of physiological and biochemical processes in plants and leads to a decrease in their productivity [6].

Based on the physiological characteristics of plants, the selection of a number of leguminous crops, among other areas, justifies the feasibility of creating genotypes with high intensity of growth processes in the early stages of development to isolate drought-resistant genotypes [3]. Studies conducted on peas have shown that within the optimal length of the stem for a specific soil-climatic zone, genotypes that differ in faster germinal root growth rates have advantages in crop formation [4]. This provides favorable starting conditions for the development of plants, *contributing to the* good rooting of seedlings, an increase in the absorbing surface of the root system, as well as drought tolerance [5]. Currently, there are many physiological methods for assessing the drought tolerance of field crops. The simplest, indirect methods for mass estimation of relative drought tolerance are based on the determination of seed germination and the growth of seedlings in osmotic solutions imitating moisture. The ability of seeds to germinate under these conditions reflects, on the one hand, the hereditary characteristic of germination with relatively less water, on the other, the presence of a high suction force, which ensures the rapid absorption of the required amount of water [8]. The high sucking power of the seeds causes not only better germination with a lack of moisture, but also the formation of a more powerful root system (primitive), which is important for the further life of the plants, especially during drought, i.e. properties of the seedling to a large extent affect the formation of drought tolerance in an adult plant [2]. In this regard, it is necessary to select the adaptive and most drought-resistant variety samples.

Methods. The objects of research were 31 collection samples of lentils from different countries. The laboratory studies on the assessment of the drought resistance of lentil plants were carried out in the laboratory "Analysis of grain quality" LLP "Kazakh Research Institute of Agriculture and Crop Production". Drought tolerance relative lentil accessions were evaluated in the early stages of plant growth by seed germination in the sand in 3 cases (PEG-10%, PEG-15%, control is distilled water) to 3-fold replicates. The seeds were germinated in a thermostat, at a temperature of +20 °C in the dark. Seed germination was taken into account on day 7. It was dried in a drying cabinet at a temperature of +130 °C, 40 minutes. According to the methodological guide All-Russian Institute of Plant Industry 1988 Udovenko G.V. [2], solutions with different concentrations of PEG 6000 were prepared according to the method of Michel B. E., Kaufmann M. R. [7].

The percentage of germinated seeds (P) is determined as follows:

The number of seeds germinated in the control is taken as 100%, the average number of seeds germinated in an osmotic stress medium (PEG 15% and PEG 10%) (a) is expressed as a percentage of the number of seeds germinated in the control (b) thus formula 1:

$$P = \frac{a}{b} * 100\% . \quad (1)$$

The higher the percentage of seed germination in the PEG nutrient medium is 15%, the more drought-resistant the sample.


To determine depression of growth processes (Index of tolerance), the average dry weight of the seedlings in the control is taken as 100% (x), the average dry weight in the osmotic stress medium (PEG 10% and PEG 15%) (y) is determined in percent of dry weight by seedlings with increased osmotic pressure (z) is determined by the formula 2:

$$Z = 100 - y/x * 100 . \quad (2)$$

In more drought tolerance, the accumulation of seedlings of biomass is inhibited to a lesser extent.

Results. In determining the germination of various levels of osmotic stress of lentils, no significant differences were found. To characterize the drought tolerance of plants, these lengths and numbers of germinal roots were used, since the ability to form a powerful primary root system plays an important role in drought conditions.

The lack of water caused by various concentrations of PEG led to a decrease in stem height, root length and seedling mass, for different lentil samples (figure).



Seedlings on day 7: a – control, b – PEG-10%, c – PEG -15%

In control, stem height and root length was longer than sprouts in an osmotic stress environment.

Table 1 – Characteristics of drought resistance at stem height and germ-length roots

Sample name	Stemheight, cm			Germ length, cm		
	Control	PEG 10%	PEG 15%	Control	PEG 10%	PEG 15%
4605	10	4.1	3.5	4.2	3.4	2.8
K-184	10	6.2	4.	4.5	3.6	2.5
LC046000246L	10	4.3	3.2	4.5	3.7	2.3
39113	9.9	4.9	3.1	3.9	3.3	2.4
LC04600023L	9.6	4.3	3.5	4.1	2.8	2.8
K-2849	9.5	4.6	3.4	4.8	3.7	2.9
LC04600017L	9.4	4.4	4.1	3.8	3.8	3.2
LC04600068L	9.2	3.9	3.2	3.3	3	2.9
23108	9.2	4.9	3.8	4.3	3.2	2.9
Vekhovskaya	9.1	3.9	2.7	5.1	3.7	3.2
LC046000213L	9.1	4.3	3.5	3.7	3.3	3
23208	8.9	4.8	3.3	4.3	3.3	1.9
39126	8,8	4.1	3.3	3.9	3.5	3.3
K-1975	8.7	4.7	3.7	3.5	2.8	2.5
LC046000103L	8.7	4.5	2.8	4.9	3.6	3.3
23209	8.7	3.9	3.2	3.3	3.3	3.2
LC046000270L	8.6	5.5	4.1	4.4	4.2	3.4
K-2017	8.5	5.2	3.6	3.7	3.1	1.9
39119	8.5	3.6	3.5	3.9	3.4	3.1
23202	8.4	4.2	3.3	3.8	3.5	2.3
LC046000150L	8.4	3.8	3.3	3.3	3.5	3
31215	8.2	4.4	4.	3.5	3.2	2.9
LC046000156L	8.2	4.1	2.4	3.5	3.1	2.7
39203	8.1	4.6	4.2	3.5	2.9	2.9
LC046000223L	8.1	4.1	4.1	3.4	3.5	3.4
LC04600010L	7.9	4.5	3.7	3.5	3.3	3
K-6	7.9	7.5	4.1	5.9	2.9	2.5
LC046000170L	7,8	3.8	3.3	3.1	3.1	2.7
LC046000202L	7.7	4.1	3.1	3.97	2.4	2.4
39227	7	3	2.6	2.7	2.6	2.4
39229	7.1	4.3	3.3	3.4	2.8	2.2

The height of the stem under the conditions of osmotic PEG-10% allocated breeding numbers (7.5-5.2 cm): K-184, LC046000270L, K-2017, K-6, with PEG-15% stood out (4.2-4 cm): K-184, LC04600017L, LC046000270L, 31215, 39203, LC046000223L, K-6.

The length of the root in the osmotic medium PEG-10% stood out (4.2-3.5 cm): LC046000270L, when PEG-15% stood out (3.4-3.1 cm): 39126, LC046000103L, LC046000270L, LC046000223L sorrow samples.

Table 2 –Evaluation of drought tolerance by green and dry weight of lentil seedlings

Sample name	Green mass, g			Dry weight, g		
	Control	PEG 10%	PEG 15%	Control	PEG 10%	PEG 15%
4605	5.8	2	1.9	1.2	0.8	0.5
K-184	1.5	1.4	1.2	0.3	0.4	0.4
LC046000246L	2.5	2	1.5	0.7	0.7	0.5
39113	2.3	1.6	1.6	1.	0.5	0.4
LC04600023L	1.8	1.6	1.6	0.7	0.6	0.4
K-2849	2.6	2.1	1.6	0.8	0.6	0.6
LC04600017L	1.7	1.5	1,3	0.6	0.5	0.3
LC04600068L	2.3	1.6	1.4	0.7	0.6	0.6
23108	1.2	1.	0.9	0.3	0.5	0.4
Vekhovskaya	4.8	1.79	1.2	1.2	0.7	0.5
LC046000213L	2.7	2.1	1,3	0.9	0.7	0.5
23208	1.87	1.5	1.8	0.5	0.5	0.6
39126	1.7	1.6	1.1	0.6	0.5	0.4
K-1975	1,3	1.29	1.2	0.6	0.2	0.4
LC046000103L	1.97	1.6	1.6	0.8	0.7	0.4
23209	2.1	1.9	1.9	0.9	0.9	0.5
LC046000270L	2.1	1.5	1.6	0.9	0.6	0.5
K-2017	2.8	2.2	1.7	0.6	0.4	0.3
39119	1.8	1.7	1.5	0.8	0.5	0.5
23202	2.4	1,3	1.9	0.7	0.6	0.6
LC046000150L	1.7	1.5	1.4	0.6	0.6	0.5
31215	1.6	1.5	1.2	0.6	0.5	0.3
LC046000156L	3.7	2.3	1.7	0.7	0.6	0.45
39203	1.4	0.9	0.9	0.5	0.4	0.2
LC046000223L	1.7	1.7	1.6	0.6	0.6	0.5
LC04600010L	1.7	1.5	1.5	0.8	0.6	0.5
K-6	5.9	5.3	2.1	1.1	0.3	0.3
LC046000170L	4.	2.54	2.1	0.9	0.6	0.9
LC046000202L	1.8	1.7	1.56	0.9	0.6	0.4
39227	7,6	3.9	1.4	0.5	0.6	0.6
39229	4.01	2.8	1.4	0.4	0.4	0.3

According to the green mass in the osmotic medium PEG 10% stood out (0.9-5.3 g): K-6, LC046000170L, 39227, with PEG-15% (0.9-2.1 g): K-6, LC046000170L variety samples.

By dry mass, promising in an osmotic medium is 10% PEG-e (0.4-1.2 g): K-6, LC046000170L, with PEG-15% (0.3-0.9 g): K-6, LC046000170L variety samples.

In more drought-resistant varieties, the accumulation of seedlings of biomass is inhibited to a lesser extent.

Table 3 – Analysis of the resistance index of lentil samples with osmotic stress PEG-10%

Name of variety samples	Origin	Sustainability Index %
		PEG 10%
Resistant variety samples		
K-6	Russia	72.7
K-1975	Canada	66.7
39113	Russia	50
Medium StableSamples		
Vekhovskaya	Russia	41.7
39119	Russia	37.5
4605	Russia	33.3
K-2017	Canada	33.3
LC046000202L	Syria	33.3
LC046000270L	Syria	33.3
K-2849	Russia	25
LC04600010L	Syria	25
LC046000213L	Syria	22.2
39203	Russia	20
39126	Russia	16.7
LC04600017L	Syria	16.7
31215	Russia	16.7
LC04600023L	Syria	14.3
23202	Russia	14.3
LC04600068L	Syria	14.3
LC046000156L	Syria	14.3
LC046000103L	Syria	12.5
Sensitive variety samples		
23208	Russia	0
LC046000246L	Syria	0
LC046000223L	Syria	0
LC046000150L	Syria	0
23209	Russia	0
39229	Russia	0
39227	Russia	-20
K-184	Russia	-33,3
23108	Russia	-66,7

The results showed that osmotic stress has a significant impact on early vegetative developmental stages; the effect depended on the degree of stress. By definition, depression of growth processes with an osmotic stress concentration of PEG of 10%, according to the tolerance index, was divided into 3 groups:

Resistant variety samples are K-6, K-1975, 39113. Mediumly resistant variety samples are Vekhovskaya, 39119, LC046000170L, 4605, K-2017, LC046000202L, LC046000270L, K-2849, LC04600010L, LC046000213L, 39203, by the index, by the text, by the text, by the text; 23202, LC04600068L, LC046000156L, LC046000103L. Sensitive is 23208, LC046000246L, LC046000223L, LC046000150L, 23209, 39229, 39227, K-184, 23108 are variety samples.

Table 4 – Analysis of the stability index of lentil samples with osmotic stress PEG-15%

Name of variety samples	Origin	Sustainability Index %
		PEG 15%
Resistant variety samples		
K-6	Russia	72.7
39113	Russia	60
39203	Russia	60
4605	Russia	58.3
Vekhovskaya	Russia	58
LC046000202L	Syria	55.5
K-2017	Canada	50
LC046000103L	Syria	50
LC04600017L	Syria	50
31215	Russia	50
Medium StableSamples		
LC046000213L	Syria	44.4
LC046000270L	Syria	44.4
23209	Russia	44.4
LC04600023L	Syria	42,8
LC04600010L	Syria	37.5
39119	Russia	37.5
LC046000156L	Syria	35.7
K-1975	Canada	33.3
39126	Russia	33.3
LC046000246L	Syria	28.6
K-2849	Russia	25
23202	Russia	14.3
LC04600068L	Syria	14.3
39229	Russia	25
LC046000150L	Syria	16, 7
LC046000223L	Syria	16.7
23202	Russia	14.3
LC04600068L	Syria	14.3
Sensitive variety samples		
LC046000170L	Syria	0
23208	Russia	-20
39227	Russia	-20
K-184	Russia	-33,3
23108	Russia	-33,3

According to the definition of depression, growth processes with an osmotic stress concentration of PEG of 15% were divided into 3 groups:

Sustainable variety samples are K-6, 39113, 39203, 4605, Vekhovskaya, LC046000202L, K-2017, LC046000103L, LC04600017L, 31215. Mediumly resistant variety samples are LC046000213L, LC046000270L, 23209, LC04600023L, LC04600010L, 39119, LC046000156L, K-1975, 39126,

LC046000246L, K-2849, 39229, LC046000150L, LC046000223L, 23202, LC04600068L. Sensitive is LC046000170L, 23208, 39227, K-184, 23108 are variety samples.

Conclusion. The results showed that osmotic stress has a significant impact on early vegetative developmental stages; the effect depended on the degree of stress.

Analysis of lentil variety samples under osmotic stress conditions (PEG 10% and PEG 15%) revealed highly drought-resistant variety samples: K-6, 39113.

These samples will be used as a starting material for breeding work on drought tolerance.

**А. Ж. Сайкенова^{1,2}, Т. Н. Нургасенов¹, М. С. Кудайбергенов²,
С. В. Дидоренко², Б. Р. Сайкенов¹**

¹Қазақ ұлттық аграрлық университет, Алматы, Қазақстан,

²Қазақ егіншілік және өсімдік шаруашылығы ғылыми зерттеу институты, Алматы облысы, Қазақстан

ЖАСЫМЫҚ СОРТУЛГІЛЕРІН ҚҰРҒАҚШЫЛЫҚҚА ТӨЗІМДІЛІГІН САЛЫСТЫРМАЛЫ БАҒАЛАУ

Аннотация. Мақалада жасымық сортүлгілерін құрғақшылыққа төзімділігін құмда ПЭГ-10%, 15% ерітінділерімен зертханалық тәжірибе арқылы анықтау. Құрғақшылыққа қатысты деректер келтірілген. Осмотикалық стресс (ПЭГ 10% және ПЭГ 15%) талдауы бойынша жасымық сортүлгілерінен құрғақшылыққа өте төзімді үлгілері анықталды: К-6, 39113 (Ресей).

Түйін сөздер: жасымық, сортүлгілері, өңгіштік, құрғақшылыққа төзімділік, осмотикалық стресс.

**А. Ж. Сайкенова^{1,2}, Т. Н. Нургасенов¹, М. С. Кудайбергенов²,
С. В. Дидоренко², Б. Р. Сайкенов¹**

¹Казахский национальный аграрный университет, Алматы, Казахстан,

²Казахский научно-исследовательский институт земледелия и растениеводства, Алматинская область, Казахстан

СРАВНИТЕЛЬНАЯ ОЦЕНКА ЗАСУХОУСТОЙЧИВОСТИ СОРТООБРАЗЦОВ ЧЕЧЕВИЦЫ

Аннотация. В статье приведены результаты лабораторных опытов по определению относительной засухоустойчивости сортообразцов чечевицы методом проращивания семян на песке в растворе ПЭГ - 10%, 15%. Приводятся данные по засухоустойчивости. Анализ сортообразцов чечевицы в условиях осмотического стресса (ПЭГ 10% и ПЭГ 15%) выявил высокозасухоустойчивые сортообразцы: К-6, 39113 (Россия).

Ключевые слова: чечевица, сортообразцы, всхожесть, засухоустойчивость, осмотический стресс.

Information about authors:

Saikenova Alma Zhumabayevna, PhD student of 3 year-study at the Kazakh National Agrarian University, Almaty, Kazakhstan; alma.arai@mail.ru; <https://orcid.org/0000-0002-9211-1415>

Nurgasenov Taken Nurgasenovich, Professor, Doctor of Agricultural Sciences, Kazakh National Agrarian University, Almaty, Kazakhstan; nurgasenov.t@mail.ru; <https://orcid.org/0000-0002-1441-9596>

Kudaibergenov Mukhtar Sarsenbekovich, Doctor of Biological Sciences, Academician ASKhN RK, Kazakh Research Institute of Agriculture and Crop Production, Almaty region, Kazakhstan; muhtar.sarsenbek@mail.ru; <https://orcid.org/0000-0002-8185-3990>

Didorenko Svetlana Vladimirovna, Candidate of Biological Sciences, Head of Leguminous Crops Department, Kazakh Research Institute of Agriculture and Crop Production, Almaty region, Kazakhstan; svetl_did@mail.ru; <https://orcid.org/0000-0002-2223-0718>

Saykenov Bakytzhan Rahmetollinovich, Associate Professor, Candidate of Agricultural Sciences, Kazakh National Agrarian University, Almaty, Kazakhstan; saikenov67@gmail.ru

REFERENCES

- [1] The market for lentils of Russia in 2011-2013, January-March 2014 [Electronic resource], access mode: <http://www.stgetman.narod.ru/checheva.html>.
- [2] Udovenko G.V. Diagnostics of plant resistance to stress: Methodical manual. VIR. Leningrad, 1988. 226 p.
- [3] Debely G.A. Leguminous crops in the Nechernozemny zone of the Russian Federation. M.: Nemchinovka, 2009. 260 p.
- [4] Novikova N.E. Study of the interrelated relations between the aerial part and the root system near the mountains in connection with the tasks of selection // Biological and economic potential of leguminous, cereal crops and the ways of its realization. Orel, 1999. P. 62-66.
- [5] Novikova N.E., Uvarov V.N., Kondykov I.V. Use in the selection of peas of a new method of selection according to the parameters of plant growth at the early stage of ontogenesis // Vestnik RAAS. 2007. N 6. P. 43-46.
- [6] Naumkin V.N., Naumkin L.A., Kurenskaya O.Yu., Artyukhov A.I., Lukashevich M.I., Ageeva P.A. Comparative assessment of the drought resistance of varieties and varieties of forage lupine // Plant. 2015. N 3. P. 10-11.
- [7] Michel B.E., Kaufmann M.R. (1973). The Osmotic Potential of Polyethylene Glycol6000 // Plant Physiology. 51: 914-916.
- [8] Dazova A.M., Dyrka S. The definition and essence of land relations as an economic category // 2018. Vol. 6, N 376. P. 67-73. ISSN 2518-1467.29 (onlain). ISSN 1991-3494 (Print). <https://doi.org/10.32014/2018.2518-1467.29>

Publication Ethics and Publication Malpractice in the journals of the National Academy of Sciences of the Republic of Kazakhstan

For information on Ethics in publishing and Ethical guidelines for journal publication see <http://www.elsevier.com/publishingethics> and <http://www.elsevier.com/journal-authors/ethics>.

Submission of an article to the National Academy of Sciences of the Republic of Kazakhstan implies that the described work has not been published previously (except in the form of an abstract or as part of a published lecture or academic thesis or as an electronic preprint, see <http://www.elsevier.com/postingpolicy>), that it is not under consideration for publication elsewhere, that its publication is approved by all authors and tacitly or explicitly by the responsible authorities where the work was carried out, and that, if accepted, it will not be published elsewhere in the same form, in English or in any other language, including electronically without the written consent of the copyright-holder. In particular, translations into English of papers already published in another language are not accepted.

No other forms of scientific misconduct are allowed, such as plagiarism, falsification, fraudulent data, incorrect interpretation of other works, incorrect citations, etc. The National Academy of Sciences of the Republic of Kazakhstan follows the Code of Conduct of the Committee on Publication Ethics (COPE), and follows the COPE Flowcharts for Resolving Cases of Suspected Misconduct (http://publicationethics.org/files/u2/New_Code.pdf). To verify originality, your article may be checked by the Cross Check originality detection service <http://www.elsevier.com/editors/plagdetect>.

The authors are obliged to participate in peer review process and be ready to provide corrections, clarifications, retractions and apologies when needed. All authors of a paper should have significantly contributed to the research.

The reviewers should provide objective judgments and should point out relevant published works which are not yet cited. Reviewed articles should be treated confidentially. The reviewers will be chosen in such a way that there is no conflict of interests with respect to the research, the authors and/or the research funders.

The editors have complete responsibility and authority to reject or accept a paper, and they will only accept a paper when reasonably certain. They will preserve anonymity of reviewers and promote publication of corrections, clarifications, retractions and apologies when needed. The acceptance of a paper automatically implies the copyright transfer to the National Academy of Sciences of the Republic of Kazakhstan.

The Editorial Board of the National Academy of Sciences of the Republic of Kazakhstan will monitor and safeguard publishing ethics.

Правила оформления статьи для публикации в журнале смотреть на сайте:

www:nauka-nanrk.kz

<http://agricultural.kz/index.php/en/>

Редактор *М. С. Ахметова, Т. М. Апендиев, Д. С. Аленов*
Верстка на компьютере *Д. Н. Калкабековой*

Подписано в печать 11.06.2019.
Формат 60x881/8. Бумага офсетная. Печать – ризограф.
5,75 п.л. Тираж 300. Заказ 3.