

ISSN 2224-526X

ҚАЗАҚСТАН РЕСПУБЛИКАСЫ  
ҰЛТТЫҚ ҒЫЛЫМ АКАДЕМИЯСЫНЫҢ  
Қазақ ұлттық аграрлық университеті

# Х А Б А Р Л А Р Ы

## ИЗВЕСТИЯ

НАЦИОНАЛЬНОЙ АКАДЕМИИ НАУК  
РЕСПУБЛИКИ КАЗАХСТАН  
Қазақстан Республикасының  
Ұлттық ғылым академиясының  
Қазақ ұлттық аграрлық университеті

## IZVESTIĀ

NATIONAL'NOJ AKADEMII NAUK  
RESPUBLIKI KAZAHSTAN  
Kazakh national  
agrarian university

SERIĀ AGRARNYH NAUK

4 (52)

JULY – AUGUST 2019

PUBLISHED SINCE JANUARY 2011

PUBLISHED 6 TIMES A YEAR

ALMATY, NAS RK

Б а с р е д а к т о р

**Есполов Т.И.,**

э.ғ.д, профессор,

ҚР ҰҒА академигі және вице-президенті

Р е д а к ц и я а л қ а с ы:

**Байзақов С.Б.**, э.ғ.д, проф., ҚР ҰҒА академигі (бас редактордың орынбасары); **Тиреуов К.М.**, э.ғ.д, проф., ҚР ҰҒА академигі (бас редактордың орынбасары); **Елешев Р.Е.**, т.ғ.д, проф., ҚР ҰҒА академигі; **Рау А.Г.**, т.ғ.д, проф., ҚР ҰҒА академигі; **Иванов Н.П.**, в.ғ.д, проф., ҚР ҰҒА академигі; **Кешуов С.А.**, т.ғ.д, проф., ҚР ҰҒА академигі; **Мелдебеков А.**, а.ш.ғ.д., проф., ҚР ҰҒА академигі; **Чоманов У.Ч.**, т.ғ.д, проф., ҚР ҰҒА академигі; **Елюбаев С.З.**, а.ш.ғ.д., проф., ҚР ҰҒА академигі; **Садықұлов Т.**, а.ш.ғ.д., проф., академигі; **Баймұқанов Д.А.**, а.ш.ғ.д., проф., ҚР ҰҒА корр-мүшесі; **Сансызбай А.Р.**, а.ш.ғ.д., проф., ҚР ҰҒА корр-мүшесі; **Умбетаев И.**, а.ш.ғ.д., проф., ҚР ҰҒА академигі; **Оспанов С.Р.**, а.ш.ғ.д., проф., ҚР ҰҒА құрметті мүшесі; **Олейченко С.И.**, а.ш.ғ.д., проф.; **Кененбаев С.Б.**, а.ш.ғ.д., проф., ҚР ҰҒА корр-мүшесі; **Омбаев А.М.**, а.ш.ғ.д., проф. ҚР ҰҒА корр-мүшесі; **Молдашев А.Б.**, э.ғ.д, проф., ҚР ҰҒА құрметті мүшесі; **Сагитов А.О.**, б.ғ.д, ҚР ҰҒА академигі; **Сапаров А.С.**, а.ш.ғ.д., проф., ҚР АШҒА академигі; **Балгабаев Н.Н.**, а.ш.ғ.д., проф.; **Умирзаков С.И.**, т.ғ.д, проф.; **Султанов А.А.**, в.ғ.д, проф., ҚР АШҒА академигі; **Алимкулов Ж.С.**, т.ғ.д, проф., ҚР АШҒА академигі; **Сарсембаева Н.Б.**, в.ғ.д, проф.

Р е д а к ц и я к е ñ е с і:

**Fasler-Kan Elizaveta**, Dr., University of Basel Switzerland; **Koolmees Petrus Adrianus**, Prof. Dr., Utrecht University, The Netherlands; **Babadoost-Kondri Mohammad**, Prof., University of Illinois, USA; **Yus Aniza Binti Yusof**, Dr., University Putra, Malaysia; **Hesseln Hayley Fawn**, As. Prof., University of Saskatchewan, Canada; **Alex Morgounov**, Pr., International Maize and Wheat Improvement Center Turkey; **Андреш С.**, Молдова Республикасы ҰҒА академигі; **Гаврилюк Н.Н.**, Украина ҰҒА академигі; **Герасимович Л.С.**, Беларусь Республикасының ҰҒА академигі; **Мамедов Г.**, Азербайжан Республикасының ҰҒА академигі; **Шейко И.П.**, Беларусь Республикасының ҰҒА академигі; **Жалнин Э.В.**, т.ғ.д., проф., Ресей; **Боинчан Б.**, а.ш.ғ.д, проф., Молдова Республикасы; **Юлдашбаев Ю.А.**, а.ш.ғ.д, проф., РФА корр-мүшесі, Ресей.

Главный редактор

**Есполов Т.И.,**

доктор эконом. наук, проф.,  
вице-президент и академик НАН РК

Редакционная коллегия:

**Байзаков С.Б.**, доктор эконом. наук, проф., академик НАН РК (заместитель главного редактора); **Тиреуов К.М.**, доктор эконом. наук, проф., академик НАН РК (заместитель главного редактора); **Елешев Р.Е.**, доктор техн. наук, проф., академик НАН РК; **Рау А.Г.**, доктор техн. наук, проф., академик НАН РК; **Иванов Н.П.**, доктор ветеринар. наук, проф., академик НАН РК; **Кешуов С.А.**, доктор техн. наук, проф., академик НАН РК; **Мелдебеков А.**, доктор сельхоз. наук, проф., академик НАН РК; **Чоманов У.Ч.**, доктор техн. наук, проф., академик НАН РК; **Елюбаев С.З.**, доктор сельхоз. наук, проф., академик НАН РК; **Садыкулов Т.**, доктор сельхоз. наук, проф., академик НАН РК; **Баймуқанов Д.А.**, доктор сельхоз. наук, проф., член-корр. НАН РК; **Сансызбай А.Р.**, доктор сельхоз. наук, проф., член-корр. НАН РК; **Умбетаев И.**, доктор сельхоз. наук, проф., академик НАН РК; **Оспанов С.Р.**, доктор сельхоз. наук, проф., Почетный член НАН РК; **Олейченко С.И.**, доктор сельхоз. наук, проф.; **Кененбаев С.Б.**, доктор сельхоз. наук, проф., член-корр. НАН РК; **Омбаев А.М.**, доктор сельхоз. наук, проф. член-корр. НАН РК.; **Молдашев А.Б.**, доктор эконом. наук, проф., Почетный член НАН РК; **Сагитов А.О.**, доктор биол. наук, академик НАН РК; **Сапаров А.С.**, доктор сельхоз. наук, проф., академик АСХН РК; **Балгабаев Н.Н.**, доктор сельхоз. наук, проф.; **Умирзаков С.И.**, доктор техн. наук, проф.; **Султанов А.А.**, доктор ветеринар. наук, проф., академик АСХН РК; **Алимкулов Ж.С.**, доктор техн. наук, проф., академик АСХН РК; **Сарсембаева Н.Б.**, доктор ветеринар. наук, проф.

Редакционный совет:

**Fasler-Kan Elizaveta**, Dr., University of Basel Switzerland; **Koolmees Petrus Adrianus**, Prof. Dr., Utrecht University, The Netherlands; **Babadoost-Kondri Mohammad**, Prof., University of Illinois, USA; **Yus Aniza Binti Yusof**, Dr., University Putra, Malaysia; **Hesseln Hayley Fawn**, As.Prof., University of Saskatchewan, Canada; **Alex Morgounov**, Pr., International Maize and Wheat Improvement Center Turkey; **Андреш С.**, академик НАН Республики Молдова; **Гаврилюк Н.Н.**, академик НАН Украины; **Герасимович Л.С.**, академик НАН Республики Беларусь; **Мамедов Г.**, академик НАН Республики Азербайджан; **Шейко И.П.**, академик НАН Республики Беларусь; **Жалнин Э.В.**, доктор техн. наук, проф., Россия; **Боинчан Б.**, доктор сельхоз. наук, проф., Республика Молдова; **Юлдашбаев Ю.А.**, доктор сельхоз. наук, проф., член-корр. РАН, Россия.

**Известия Национальной академии наук Республики Казахстан. Серия аграрных наук.**

**ISSN 2224-526X**

Собственник: ООО «Национальная академия наук Республики Казахстан» (г. Алматы)

Свидетельство о постановке на учет периодического печатного издания в Комитете информации и архивов Министерства культуры и информации Республики Казахстан № 10895-Ж, выданное 30.04.2010 г.

Периодичность 6 раз в год

Тираж: 300 экземпляров

Адрес редакции: 050010, г. Алматы, ул. Шевченко, 28, ком. 219-220, тел. 272-13-19, 272-13-18

<http://agricultural.kz/index.php/en/>

---

© Национальная академия наук Республики Казахстан, 2019

Адрес типографии: ИП «Аруна», г. Алматы, ул. Муратбаева, 75

Chief Editor

**Espolov T.I.,**

Dr. economy. Sciences, prof.,  
Vice President and academician of the NAS RK

Editorial Board:

**Baizakov S.B.**, Dr. of economy sciences, prof., academician of NAS RK (deputy editor); **Tireuov K.M.**, Doctor of Economy Sciences., prof., academician of NAS RK (deputy editor); **Eleshev R.E.**, Dr. Of agricultural sciences, prof., academician of NAS RK; **Rau A.G.**, Dr. sciences, prof., academician of NAS RK; **Ivanov N.P.**, Dr. of veterinary sciences, prof., academician of NAS RK; **Keshuov S.A.**, Dr. sciences, prof., academician of NAS RK; **Meldebekov A.**, doctor of agricultural sciences, prof., academician of NAS RK; **Chomanov U.Ch.**, Dr. sciences, prof., academician of NAS RK; **Yelyubayev S.Z.**, Dr. of agricultural sciences, prof., academician of NAS RK; **Sadykulov T.**, Dr. Farm. Sciences, prof., academician of NAS RK; **Baimukanov D.A.**, doctor of agricultural sciences, prof., corresponding member NAS RK; **Sansyzbai A.R.**, doctor of agricultural sciences, prof., corresponding member NAS RK; **Umbetaev I.**, Dr. Farm. Sciences, prof., academician of NAS RK; **Ospanov S.R.**, Dr. agricultural sciences, prof., Honorary Member of NAS RK; **Oleychenko S.N.**, Dr. Of agricultural sciences, prof.; **Kenenbayev S.B.**, Dr. Agricultural sciences, prof., corresponding member NAS RK; **Ombayev A.M.**, Dr. Agricultural sciences, Prof. corresponding member NAS RK; **Moldashev A.B.**, Doctor of Economy sciences, prof., Honorary Member of NAS RK; **Sagitov A.O.**, Dr. biol. sciences, academician of NAS RK; **Saparov A.S.**, Doctor of agricultural sciences, prof., academician of NAS RK; **Balgabaev N.N.**, the doctor agricultural sciences, Prof.; **Umirzakov S.I.**, Dr. Sci. Sciences, Prof.; **Sultanov A.A.**, Dr. of veterinary sciences, prof., academician of the Academy of Agricultural Sciences of Kazakhstan; **Alimkulov J.C.**, Dr. of tekhncial sciences, prof., academician of the Academy of Agricultural sciences of Kazakhstan; **Sarsembayeva N.B.**, Dr. veterinary sciences, prof.

Editorial Board:

**Fasler-Kan Elizaveta**, Dr., University of Basel Switzzeland; **Koolmees Petrus Adrianus**, Prof. Dr., Utrecht University, The Netherlands; **Babadoost-Kondri Mohammad**, Prof., University of Illinois, USA; **Yus Aniza Binti Yusof**, Dr., University Putra, Malaysia; **Hesseln Hayley Fawn**, As. Prof., University of Saskatchewan, Canada; **Alex Morgounov**, candidate of agricultural sciences, International Maize and Wheat Improvement Center Turkey; **Andresh S.**, academician of NAS of Moldova; **Gavriluk N.N.**, academician of NAS of Ukraine; **Gerasimovich L.S.**, academician of NAS of Belorassia; **Mamadov G.**, academician of NAS of Azerbaijan; **Sheiko I.P.**, academician of NAS of Belorassia; **Zhalnin E.V.**, Dr. of technical sciences, professor, Russia, **Boinchan B.**, doctor of agricultural sciences, prof., Moldova; **Yuldashbayev Y.A.**, doctor of agricultural sciences, prof., corresponding member of RAS, Russia.

**News of the National Academy of Sciences of the Republic of Kazakhstan. Series of Agrarian Sciences.**

**ISSN 2224-526X**

Owner: RPA "National Academy of Sciences of the Republic of Kazakhstan" (Almaty)

The certificate of registration of a periodic printed publication in the Committee of Information and Archives of the Ministry of Culture and Information of the Republic of Kazakhstan N 10895-Ж, issued 30.04.2010

Periodicity: 6 times a year

Circulation: 300 copies

Editorial address: 28, Shevchenko str., of.219-220, Almaty, 050010, tel. 272-13-19, 272-13-18,  
<http://nauka-nanrk.kz/agricultural.kz>

---

© National Academy of Sciences of the Republic of Kazakhstan, 2019

Address of printing house: ST "Aruna", 75, Muratbayev str, Almaty

## NEWS

OF THE NATIONAL ACADEMY OF SCIENCES OF THE REPUBLIC OF KAZAKHSTAN

SERIES OF AGRICULTURAL SCIENCES

ISSN 2224-526X

Volume 4, Number 52 (2019), 5 – 10

<https://doi.org/10.32014/2019.2224-526X.41>

UDC 631.42

**M. R. Kerimkulova<sup>1,2</sup>, F. E. Kozybaeva<sup>3</sup>,  
Z. A. Mansurov<sup>2</sup>, Zh. O. Oshakbayeva<sup>1</sup>, A. R. Kerimkulova<sup>2</sup>**

<sup>1</sup>Kazakh national agrarian university, Almaty, Kazakhstan,

<sup>2</sup>Institute of Combustion problems, Almaty, Kazakhstan,

<sup>3</sup>Kazakh research institute of soil science and agrochemistry after U. U. Usmanov, Almaty, Kazakhstan.

E-mail: biomak111@mail.ru

## EFFECTS OF BIOCHAR ON THE POLLUTED HEAVY METALS LEACHED CHERNOZEM SOIL

**Abstract.** In a 40-day laboratory experiment, a study was conducted of the effect of biochar on the properties of leached chernozem soil contaminated with heavy metals. Biochar exhibits a great potential to efficiently tackle soil contaminants considering the wide availability of feedstock, low-cost and favorable physical/chemical surface characteristics. The well-distributed pore network that includes micropores <2 nm, mesopores 2-50 nm and macropores >50 nm. It is established that the introduction of biochar led to an increase in NPK, a decrease in the content of heavy metals Zn and Pb in the soil. It is shown that the introduction of 2% biochar positively affects the growth of vegetation and seed germination in quantity increases 31 to 58 seeds.

**Keywords:** biochar, leached chernozem, humus, mobile phosphorus, total nitrogen, exchange potassium.

**Introduction.** The presence of excessive metal contents in soils may have serious consequences for surrounding ecosystems, groundwater, agricultural productivity and human health [1]. The emergence of new concepts and requirements in the sustainable development of the world economy could not but affect such an important sector as agriculture, which directly depends on nature and causes no small environmental damage [2]. The restoration of hazardous soils is thus essential. Conventional remediation techniques (e.g. excavation, landfilling, soil washing) are nowadays recognized as inappropriate because they generate considerable disturbance in the environment and they are economically unfeasible on a large scale. The restoration of hazardous soils is thus essential. Conventional remediation techniques (e.g. excavation, landfilling, soil washing) are nowadays recognized as inappropriate because they generate considerable disturbance in the environment and they are economically unfeasible on a large scale. An alternative technique consists in introducing in situ amendments into the contaminated soils in order to reduce the metal mobility while simultaneously creating conditions that promote plant growth. This technique has received a growing interest and is turning out to be a promising green and cost-effective alternative for soil remediation [3-6]. Thus, the natural growth of plants in technogenic landscapes, regardless of the climatic zone, the type of damage and the types of discoveries, is accompanied by slow growth and the plant cannot regenerate naturally [7].

Among the methods, biosorption technique is the most common and cost-effective. This is because biosorbents are environmentally friendly and readily available in large quantities, and one of the most popular biosorbents is biochar. Biochar is a carbon-rich, fine-grained, and porous material. It has received increasing attention due to its ability to store large amount of carbon, increase crop yield, reduce soil emission of greenhouse gases, improve soil quality, decrease nutrient leaching, and reduce irrigation and fertilizer requirements [8-10].

As far as the former application is concerned, biochar has demonstrated to be able to improve soil properties [11, 12] as well as to reduce contaminant leaching [13, 14]. As adsorbent, it represents a more

economically and environmentally sustainable alternative to commercial media because its use allows to avoid industrial activities for adsorbent production as well as to reuse a waste that alternatively would need to be disposed of. As adsorbent, biochar can be applied to water and wastewater treatment, for a wide range of pollutants such as lead arsenic, copper, cadmium, chromium, mercury, zinc and nickel [15-17]. Biochar, which can be applied to contaminated soils to reduce metals mobilization. Indeed, heavy metals become bound to carbonates and organic matter after biochar incorporation, which enhances the adsorption process due to metals building bonds with oxygen, carbon and nitrogen containing functional groups [18-20].

Quality and properties of biochar depend on many factors [21]; it is essential to identify characteristics of this adsorbent to predict its effectiveness towards the target contaminant. Many biochars have a high adsorption capacity for metal contaminants because of the high heterogeneous specific surface [22] and the well-distributed pore network that includes micropores (<2 nm), mesopores (2-50 nm) and macropores (> 50 nm) [23].

**Material and methods.** *Soil sampling and characterization.* The objects of study was leached black soil contaminated with heavy metals from emissions from the zinc plant of the Ridder city, East Kazakhstan region. Soil samples were collected from 0 to 30 cm depth [24]. When dry samples were homogenised and sieved (<2 mm). In soil samples, total nitrogen, mobile phosphorus ( $P_2O_5$ ) and exchange potassium ( $K_2O$ ) [25], the pH of the aqueous extract [26], the humus content according to generally accepted methods [27], Zn and Pb by atomic absorption spectroscopy were determined [28].

*Production and characterization of biochar.* Biochar was produced from rice husk which is an abundantly available agri-food residue in Kazakhstan. Then it was placed in a reactor to prevent  $O_2$  input. The reactor furnace was heated at a rate of  $50\text{ cm}^3/\text{min}$  up to  $650^\circ\text{C}$ . The temperature was held for about one hour and subsequently cooled slowly to room temperature.

**Results and discussion.** The results of polluted soil sample analyses: pH-7.85; humus content 5.31%; total nitrogen 0,120%; mobile phosphorus ( $P_2O_5$ ) 35 mg/kg; exchange ( $K_2O$ ) 433.8 mg/kg; zinc 8804.84 mg/kg; lead 429.2 mg/kg. Our data shows that heavy metals Pb and Zn exceed the MPC.

To obtain biochar, rice husk was used (Bakanas, Almaty region), which is a renewable waste of plant origin. In order to extract the powdered remains of rice, the husk was washed with water and dried in a drying cabinet at a temperature of  $120^\circ\text{C}$ . Next, the husk was carbonized in a laboratory reactor [28] at a temperature of  $650^\circ\text{C}$  for one hour in an inert atmosphere of Ar flow. Quality and properties of biochar depend on many factors [15] it is essential to identify characteristics of this adsorbent to predict its effectiveness towards the target contaminant. We studied biochar with a particle size of not more than 1 cm, obtained from rice husk. The resulting biochar had the following properties: humidity - 1.7%, carbon - 39.98%, oxygen - 34.53%, silicon - 25.49%. In the experiment used the original soil and soil, which made biochar. Biochar properties vary considerably with feedstock material and pyrolysis temperature, with high temperature producing biochars with higher surface area, porosity, mineral contents, but less functional groups. To determine the effectiveness of the biochar obtained by us, 0.5%, 1%, and 2% of biochar were added to the soil; also a control sample of soil without biochar was prepared. There were 3 replicates for each group. Samples were kept under static conditions at room temperature ( $21\pm 10^\circ\text{C}$ ). After a 40-day incubation period in the soil of biochar, the chemical and physical properties of the soil changed: humus 5.31% decreased to 4.20%. With the introduction of 2% biochar after a 40-day incubation period in soil pH 7.85 to 6.56, the content of exchangeable potassium ( $K_2O$ ) increases significantly - 433.8 mg/kg to 589.8 mg/kg, mobile phosphorus ( $P_2O_5$ ) - 35 mg/kg to 48.40 mg/kg, nitrogen 0.120% to 0.130% increased by comparison in the control variants of the experiment. These changes are due to the adsorption capacity of biochar, as well as a decrease in the alkalinity of the studied soil, which has led to an increase in the availability of substances for plants. This indicates a long-term positive effect of the use of biochar. And the content of zinc heavy metals is from 8804.84 mg/kg to 6000,5 mg/kg and lead from 429.2 mg/kg to 129.3 mg/kg.

Additionally, biomass and growth were determined, the time of death of «Aray» plant wheat plants on leached chernozem soil contaminated with heavy metals. Seeds were planted vessels, each of which contained 300 g of soil with different content of biochar - 0.5%, 1%, 2%. Soil without biochar was used as a control. Wheat seeds before sowing were calibrated, and defective ones were removed. Sowing was carried out in a moist soil at the rate of 80 seeds per vessel. None mineral or organic fertilizer was applied.

In controlled conditions: the temperature is 23-25 °C, the light period is 14/10 h (day\night) and the soil humidity is 50% of the total moisture capacity of the soil. 14 days after sowing, wheat shoots are observed.

After 25 days on the studied vessels, the grown wheat differed in the amount of each other. As can be seen in Figure 1, out of 80 wheat seeds in the control soil, 31 wheat seeds grew, and 58% of wheat seeds grew in 2% of biochar. This suggests that biochar affects vegetation growth (figure 1).




Figure 1 – The effect of biochar for wheat seedlings

The average length of the roots of wheat on the 25<sup>th</sup> day of the experiment was 1.5 cm for the control variant. The introduction of biochar into the soil led to a significant ( $p < 0.05$ ) increase in the length of wheat roots by an average of 2.6 times as compared with the control variant (figure 2).




Figure 2 – The average length of the roots of wheat

**Conclusion.** The results of the study showed that the introduction of biochar led to an increase in the mobile phosphorus ( $P_2O_5$ ) - 35 mg/kg to 48.40%, in soil pH 7.85 to 6.56, the content of exchangeable potassium ( $K_2O$ ) increases significantly - 433.8 mg/kg to 589.8 mg/kg, nitrogen 0.120% to 0.130% increased by comparison in the control variants of the experiment. The content of heavy metals in the soil

after the introduction of biochar of zinc reduces from 8804.84 mg/kg to 6000.5 mg/kg, and lead - from 429.2 mg/kg to 129.3 mg/kg. Adding 2% biochar to the soil shows that it improves wheat growth. In the study of the germination of wheat seeds with 2% biochar increased the number from 31 to 58 seeds. The introduction of biochar into the soil led to a significant ( $p < 0.05$ ) increase in the length of wheat roots by an average of 2.6 times as compared with the control variant.

**М. Р. Керімқұлова<sup>1,2</sup>, Ф. Е. Қозыбаева<sup>3</sup>,  
З. А. Мансұров<sup>2</sup>, Ж. О. Ошақбаева<sup>1</sup>, А. Р. Керімқұлова<sup>2</sup>**

<sup>1</sup>Қазақ ұлттық аграрлық университет, Алматы, Қазақстан,

<sup>2</sup>Жану проблемалар институты, Алматы, Қазақстан,

<sup>3</sup>Ө. О. Оспанов атындағы Қазақ топырақтану және агрохимия ғылыми зерттеу институты,  
Алматы, Қазақстан

### **АУЫР МЕТАЛДАРМЕН ЛАСТАНҒАН СІЛТІСІЗДЕНГЕН ҚАРА ТОПЫРАҚҚА БИОКӨМІРДІҢ ӘСЕРІ**

**Аннотация.** 40 күндік лабораториялық экспериментте ауыр металдармен ластанған сілтісізденген қара топырақтың қасиетіне биокөмірдің әсері. Биокөмір шикізатының қол жетімділігін ескере отырып, шығыны төмен, беттік ауданының физика-химиялық қасиетінің қолайлы болуы себебінен топырақты ластаушы заттармен эффективті күресуде үлкен мүмкіндікке ие. Кеуекті торларының жақсы жетілгендігі, микрокеуек  $< 2$  нм, мезокеуекті 2-50 нм және макрокеуекті  $> 50$  нм. Биокөмірді топыраққа қосқанда НРК жоғарылап, ауыр металдар Zn және Pb төмендеуі анықталды. 2% биокөмірді топыраққа қосқанда өсімдіктердің өсуіне оңтайлы әсер етіп және тұқымның өңгіштік саны 31-ден 58 тұқымға дейін артты.

**Түйін сөздер:** биокөмір, сілтісізденген қара топырақ, гумус, жылжымалы фосфор, жалпы азот, алмаспалы калий.

**М. Р. Керимкулова<sup>1,2</sup>, Ф. Е. Козыбаева<sup>3</sup>,  
З. А. Мансуров<sup>2</sup>, Ж. О. Ошакбаева<sup>1</sup>, А. Р. Керимкулова<sup>2</sup>**

<sup>1</sup>Казахский национальный аграрный университет, Алматы, Казахстан,

<sup>2</sup>Институт Проблем горения, Алматы, Казахстан,

<sup>3</sup>Казахский научно-исследовательский институт почвоведения и агрохимии им. У. У. Успанова,  
Алматы, Казахстан

### **ВЛИЯНИЕ БИОУГЛЯ НА ЗАГРЯЗНЕННОЙ ТЯЖЕЛЫМИ МЕТАЛЛАМИ ЧЕРНОЗЕМ ВЫЩЕЛОЧЕННОЙ ПОЧВЫ**

**Аннотация.** В 40-суточном лабораторном эксперименте провели исследование влияние биоугля на свойства чернозем выщелоченной почвы, загрязненного тяжелыми металлами. Биоуголь обладает огромным потенциалом для эффективной борьбы с загрязнителями почвы, учитывая широкую доступность сырья, низкую стоимость и поверхность благоприятные физико-химические характеристики. Хорошо распределенная сеть пор, которая включает микропоры  $< 2$  нм, мезопоры 2-50 нм и макропоры  $> 50$  нм. Установлено, что внесение биоугля привело увеличению НРК, снижение тяжелых металлов Zn и Pb в почве. Показано, что внесение 2% биоугля положительно влияет на рост растительности и всхожесть семян по количеству увеличивается 31 до 58 семян.

**Ключевые слова:** биоуголь, чернозем выщелоченный, гумус, подвижный фосфор, общий азот, обменный калий.

#### **Information about authors:**

Kerimkulova M. R., PhD student KazNAU, research associate, RSE «The Institute of Combustion Problems», laboratory of carbon nanomaterials and nanobiotechnology, Almaty, Kazakhstan; biomak111@mail.ru; <https://orcid.org/0000-0002-5951-9160>



Kozybaeva F. E., doctor of biological sciences, professor of the Department of Soil Ecology, Soil Science and Agrochemistry after U. U. Usmanov, Almaty, Kazakhstan; farida\_kozybaeva@mail.ru; <https://orcid.org/0000-0002-3576-548X>

Mansurov Z. A., academician of NANS of RK, professor, doctor of chemical sciences, director RSE «The Institute of Combustion Problems», Almaty, Kazakhstan; zmansurov@kaznu.kz; <https://orcid.org/0000-0002-8956-216X>

Oshakbayeva Zh. O., candidate of biological sciences, associate professor of soil science and agrochemistry, KazNAU, Almaty, Kazakhstan; zh.oryntaevna@gmail.ru; <https://orcid.org/0000-0002-4409-7444>

Kerimkulova A. R., candidate of chemical sciences, RSE «The Institute of Combustion Problems», Almaty, Kazakhstan; Head of the laboratory of carbon nanomaterials and nanobiotechnology; almusha\_84@mail.ru; <https://orcid.org/0000-0003-4660-0944>

## REFERENCES

[1] Adriano D.C. (2001) Trace elements in terrestrial environments: biogeochemistry, bioavailability, and risks of metals, second edition. Springer-Verlag, New York. ISBN-13: 978-0387986784 (in Eng.).

[2] Bastaubayeva Sh.O., Bekbatyrov M.B., Ospanova S.O. (2018) The biological efficiency of crop rotations and green manures to improve the productivity of irrigated lands // News of National academy of sciences of the Republic of Kazakhstan. Series of agricultural sciences. Vol. 5(47). P. 60-64. ISSN 2224-526. <https://doi.org/10.32014/2018.2224-526X.8>

[3] Adriano D.C., Wenzel W.W., Vangronsveld J., Bolan N.S. (2004) Role of assisted natural remediation in environmental cleanup. *Geoderma* 122:121-142. <https://dx.doi.org/10.1016/j.geoderma.2004.01.003> (in Eng.).

[4] Kumpiene J. (2010) Trace element immobilization in soil using amendments. In: Hooda, P.S. (Ed.), *Trace Element in Soils*. Wiley-Blackwell, Chichester, UK. ISBN: 978-1-405-16037-7 (in Eng.).

[5] Bolan N.S., Park J.H., Robinson B., Naidu R., Huh K.Y. (2011) Phytostabilization: A green approach to contaminant containment // In: Donald, L.S. (Ed.), *Adv. Academic Press, Agron.* P. 145-204 (in Eng.).

[6] Houben D., Pircar J., Sonnet P. (2012) Heavy metal immobilization by cost-effective amendments in a contaminated soil: Effects on metal leaching and phytoavailability // *J. Geochem. Explor.* 123:87-94. <https://doi.org/10.1016/j.gexplo.2011.10.004> (in Eng.).

[7] Kozybaeva F.E., Beiseyeva G.B., Toktar M. (2016) Assessment of level of restoration of the vegetable cover on the rekulvivation dump of the phosphorite deposit of Kokdzhon // News of National academy of sciences of the Republic of Kazakhstan. Series of agricultural sciences. Vol. 4(34). P. 68-71. ISSN 1991-3494 2224-526. <https://doi.org/10.32014/2018.2224-526X>

[8] Kimetu J.M., Lehmann J., Solomon O.N., Mugendi D.N., Kinyangi J.M., Riha S., Verchot L., Recha J.W., Pell A.N. (2008) Reversibility of soil productivity decline with organic matter of differing quality along a degradation gradient // *Ecosystems*. 11, 726-739. <https://doi.org/10.1007/s10021-008-9154-z> (in Eng.).

[9] Lehmann J. (2007) Bio-energy in the black // *Front. Ecol. Environ.* 5, 381-387. [https://doi.org/10.1890/1540-9295\(2007\)5\[381:BITB\]2.0.CO;2](https://doi.org/10.1890/1540-9295(2007)5[381:BITB]2.0.CO;2) (in Eng.).

[10] Nguyen B., Lehmann J., Kinyangi J., Smernik R., Riha S., Engelhard M. (2009) Longterm black carbon dynamics in cultivated soil // *Biogeochemistry*. 89, 163-176.

[11] Agegnehu G., Srivastava A.K., Bird M.I. (2017) The role of biochar and biochar-compost in improving soil quality and crop performance: A review // *Appl. Soil Ecol.* 119:156-170.

<https://doi.org/10.1016/j.apsoil.2017.06.008> (in Eng.).

[12] Blanco-Canqui, H. (2017) Biochar and Soil Physical Properties // *Soil Sci. Soc. Am. J.* 84:687-711. <https://doi.org/10.2136/sssaj2017.01.0017> (in Eng.).

[13] Zong Y., Wang Y., Sheng Y., Wu C., Lu S. (2017) Ameliorating soil acidity and physical properties of two contrasting texture Ultisols with wastewater sludge biochar // *Environ. Sci. Pollut. Res.*, 1-8. <https://doi.org/10.1007/s11356-017-9509-0> (in Eng.).

[14] Tang J., Zhu W., Kookana R., Katayama A. (2013) Characteristics of biochar and its application in remediation of contaminated soil // *J. Biosci. Bioeng.* 116:653-659. <https://doi.org/10.1016/j.jbiosc.2013.05.035> (in Eng.).

[15] De Gisi S., Lofrano G., Grassi M., Notarnicola M. (2016) Characteristics and adsorption capacities of low-cost sorbents for wastewater treatment: A review // *Sustain. Mater. Technol.* 9:10-40. <https://doi.org/10.1016/j.susmat.2016.06.002> (in Eng.).

[16] Ifthikar J., Wang T., Khan A., Jawad A., Sun T., Jiao X., Chen Z., Wang J., Wang Q., Wang H., Jawad A. (2017) Highly Efficient Lead Distribution by Magnetic Sewage Sludge Biochar: Sorption Mechanisms and Bench Applications // *Bioresour. Technol.* 238:399-406. <https://doi.org/10.1016/j.biortech.2017.03.133> (in Eng.).

[17] Lu H., Zhang W., Yang Y., Huang X., Wang S., Qiu R. (2012) Relative distribution of Pb<sup>2+</sup> sorption mechanisms by sludge-derived biochar // *Water Res.* 46:854-862. <https://doi.org/10.1016/j.watres.2011.11.058> (in Eng.).

[18] Yin D., Wang X., Chen C., Peng B., Tan C., Li H. (2016) Varying effect of biochar on Cd, Pb and As mobility in a multi-metal contaminated paddy soil // *Chemosphere*. 152:196-206. DOI:10.1016/j.chemosphere.2016.01.044 (in Eng.).

[19] Bashir S., Hussain Q., Akmal M., Riaz M., Hu H., Ijaz S.S., Iqbal M., Abro S., Mehmood S., Ahmad M. (2018) Sugarcane bagasse-derived biochar reduces the cadmium and chromium bioavailability to mash bean and enhances the microbial activity in contaminated soil // *J. Soils Sediments*. 18:874–886. <https://doi.org/10.1007/s11368-017-1796-z> (in Eng.).

[20] Park J.H., Lamb D., Paneerselvam P., Choppala G., Bolan N., Chung J.W. (2011) Role of organic amendments on enhanced bioremediation of heavy metal (loid) contaminated soils // *J. Hazard. Mater.* 185:549-574. DOI:10.1016/j.jhazmat.2010.09.082 (in Eng.).

[21] Tang J., Zhu W., Kookana R., Katayama A. (2013) Characteristics of biochar and its application in remediation of contaminated soil // *J. Biosci. Bioeng.* 116:653-659. <https://doi.org/10.1016/j.jbiosc.2013.05.035> (in Eng.).

[22] Kasozi G.N., Zimmerman A.R., Nkedi-Kizza P., Gao B. (2010) Catechol and Humic Acid Sorption onto a Range of Laboratory Produced Black Carbons (Biochars) // *Environ. Sci. Technol.* 44:6189-6195. DOI:10.1021/es1014423 (in Eng.).

[23] Mukherjee A., Zimmerman A.R., Harris W. (2011) Surface chemistry variations among a series of laboratory-produced biochars // *Geoderma*. 163:247-255. <https://doi.org/10.1016/j.geoderma.2011.04.021> (in Eng.).

[24] GOST 17.4.3.01-83 Nature Conservation. Soils. General requirements for sampling [Electronic resource] / Electronic text of the document prepared by CJSC "Code" and verified by: official publication Nature Protection. Soils: Sat. Gostov. M.: Standartinform, 2008. Access Mode: <http://docs.cntd.ru/document/gost-17-4-3-01-12> (in Rus.).

[25] GOST 26483-85 Preparation of salt extract and determination of its pH according to the method of TsINA O [Electronic resource] / The electronic text of the document is verified by: official publication // Coll.: Standards publishing house. 1985. Access mode: <http://docs.cntd.ru/document/gost-26483-85> (in Rus.).

[26] GOST 26205-91. Determination of mobile compounds of phosphorus and potassium according to the method of Machigin in the modification of TsINA O [Electronic resource] / Text of the document is verified by: official publication M.: Publishing house of standards, 1992. <http://docs.cntd.ru/document/gost-26205-91> (in Rus.).

[27] GOST 26213-91. Soils. Methods for the determination of organic matter [Electronic resource] / The text of the document is verified by: official publication M.: Standards Publishing, 1992. <http://docs.cntd.ru/document/gost-26213-91> (in Rus.).

[28] ST RK ISO 11047-2008. The quality of the soil. Determination of cadmium, chromium, cobalt, copper, lead, manganese, nickel and zinc in soil extracts in aqua regia. Spectrometric methods for atomic absorption in flame and electrothermal spraying (in Rus.).

[29] Azat S., Mansurov Z., Kozybaeva F., Beiseyeva G., Murataliev A. (2017) Land cover pollution from mining and metallurgical enterprises and biotechnological method of remediation using biochar // *Nanomedicine and nanoscience research*. 4:1221-7 (in Eng.).

## **Publication Ethics and Publication Malpractice in the journals of the National Academy of Sciences of the Republic of Kazakhstan**

For information on Ethics in publishing and Ethical guidelines for journal publication see <http://www.elsevier.com/publishingethics> and <http://www.elsevier.com/journal-authors/ethics>.

Submission of an article to the National Academy of Sciences of the Republic of Kazakhstan implies that the described work has not been published previously (except in the form of an abstract or as part of a published lecture or academic thesis or as an electronic preprint, see <http://www.elsevier.com/postingpolicy>), that it is not under consideration for publication elsewhere, that its publication is approved by all authors and tacitly or explicitly by the responsible authorities where the work was carried out, and that, if accepted, it will not be published elsewhere in the same form, in English or in any other language, including electronically without the written consent of the copyright-holder. In particular, translations into English of papers already published in another language are not accepted.

No other forms of scientific misconduct are allowed, such as plagiarism, falsification, fraudulent data, incorrect interpretation of other works, incorrect citations, etc. The National Academy of Sciences of the Republic of Kazakhstan follows the Code of Conduct of the Committee on Publication Ethics (COPE), and follows the COPE Flowcharts for Resolving Cases of Suspected Misconduct ([http://publicationethics.org/files/u2/New\\_Code.pdf](http://publicationethics.org/files/u2/New_Code.pdf)). To verify originality, your article may be checked by the Cross Check originality detection service <http://www.elsevier.com/editors/plagdetect>.

The authors are obliged to participate in peer review process and be ready to provide corrections, clarifications, retractions and apologies when needed. All authors of a paper should have significantly contributed to the research.

The reviewers should provide objective judgments and should point out relevant published works which are not yet cited. Reviewed articles should be treated confidentially. The reviewers will be chosen in such a way that there is no conflict of interests with respect to the research, the authors and/or the research funders.

The editors have complete responsibility and authority to reject or accept a paper, and they will only accept a paper when reasonably certain. They will preserve anonymity of reviewers and promote publication of corrections, clarifications, retractions and apologies when needed. The acceptance of a paper automatically implies the copyright transfer to the National Academy of Sciences of the Republic of Kazakhstan.

The Editorial Board of the National Academy of Sciences of the Republic of Kazakhstan will monitor and safeguard publishing ethics.

Правила оформления статьи для публикации в журнале смотреть на сайте:

**www:nauka-nanrk.kz**

<http://agricultural.kz/index.php/en/>

Редактор *М. С. Ахметова, Т. М. Апендиев, Д. С. Аленов*  
Верстка на компьютере *Д. Н. Калкабековой*

Подписано в печать 25.07.2019.

Формат 60x881/8. Бумага офсетная. Печать – ризограф.  
4,5 п.л. Тираж 300. Заказ 4.