

ISSN 2224-526X

ҚАЗАҚСТАН РЕСПУБЛИКАСЫ
ҰЛТТЫҚ ФЫЛЫМ АКАДЕМИЯСЫНЫҢ
Қазақ ұлттық аграрлық университеті

Х А Б А Р Л А Р Ы

ИЗВЕСТИЯ

НАЦИОНАЛЬНОЙ АКАДЕМИИ НАУК
РЕСПУБЛИКИ КАЗАХСТАН
Казахский национальный
аграрный университет

IZVESTIÂ

NATIONAL'NOJ AKADEMII NAUK
RESPUBLIKI KAZAHSTAN
Kazakh national
agrarian university

SERIÂ AGRARNYH NAUK

5 (53)

SEPTEMBER – OCTOBER 2019

PUBLISHED SINCE JANUARY 2011

PUBLISHED 6 TIMES A YEAR

ALMATY, NAS RK

Б а с р е д а к т о р

Есполов Т.И.,
э.ғ.д, профессор,
ҚР ҰҒА академигі және вице-президенті

Р е д а к ц и я алқасы:

Байзаков С.Б., э.ғ.д, проф., ҚР ҰҒА академигі (бас редактордың орынбасары); **Тиреуов К.М.**, э.ғ.д, проф., ҚР ҰҒА академигі (бас редактордың орынбасары); **Елешев Р.Е.**, т.ғ.д., проф., ҚР ҰҒА академигі; **Рау А.Г.**, т.ғ.д., проф., ҚР ҰҒА академигі; **Иванов Н.П.**, в.ғ.д, проф., ҚР ҰҒА академигі; **Кешуов С.А.**, т.ғ.д., проф., ҚР ҰҒА академигі; **Мелдебеков А.**, а.ш.ғ.д., проф., ҚР ҰҒА академигі; **Чоманов У.Ч.**, т.ғ.д., проф., ҚР ҰҒА академигі; **Елюбаев С.З.**, а.ш.ғ.д., проф., ҚР ҰҒА академигі; **Садыкулов Т.**, а.ш.ғ.д., проф., академигі; **Баймұқанов Д.А.**, а.ш.ғ.д., проф., ҚР ҰҒА корр-мүшесі; **Сансызбай А.Р.**, а.ш.ғ.д., проф., ҚР ҰҒА корр-мүшесі; **Умбетаев И.**, а.ш.ғ.д., проф., ҚР ҰҒА академигі; **Оспанов С.Р.**, а.ш.ғ.д., проф., ҚР ҰҒА құрметті мүшесі; **Олейченко С.И.**, а.ш.ғ.д., проф.; **Кененбаев С.Б.**, а.ш.ғ.д., проф., ҚР ҰҒА корр-мүшесі; **Омбаев А.М.**, а.ш.ғ.д., проф. ҚР ҰҒА корр-мүшесі; **Молдашев А.Б.**, э.ғ.д., проф., ҚР ҰҒА құрметті мүшесі; **Сагитов А.О.**, б.ғ.д., ҚР ҰҒА академигі; **Сапаров А.С.**, а.ш.ғ.д., проф., ҚР АШҒА академигі; **Балгабаев Н.Н.**, а.ш.ғ.д., проф.; **Умирзаков С.И.**, т.ғ.д, проф.; **Султанов А.А.**, в.ғ.д., проф., ҚР АШҒА академигі; **Алимкулов Ж.С.**, т.ғ.д., проф., ҚР АШҒА академигі; **Сарсембаева Н.Б.**, в.ғ.д., проф.

Р е д а к ц и я к е н е сі:

Fasler-Kan Elizaveta, Dr., University of Basel Switzerland; **Koolmees Petrus Adrianus**, Prof. Dr., Utrecht University, The Netherlands; **Babadoost-Kondri Mohammad**, Prof., University of Illinois, USA; **Yus Aniza Binti Yusof**, Dr., University Putra, Malaysia; **Hesseln Hayley Fawn**, As. Prof., University of Saskatchewan, Canada; **Alex Morgounov**, Pr., International Maize and Wheat Improvement Center Turkey; **Андреш С.**, Молдова Республикасы ҰҒА академигі; **Гаврилюк Н.Н.**, Украина ҰҒА академигі; **Герасимович Л.С.**, Беларусь Республикасының ҰҒА академигі; **Мамедов Г.**, Азербайджан Республикасының ҰҒА академигі; **Шейко И.П.**, Беларусь Республикасының ҰҒА академигі; **Жалнин Э.В.**, т.ғ.д., проф., Ресей; **Боинчан Б.**, а.ш.ғ.д, проф., Молдова Республикасы; **Юлдашбаев Ю.А.**, а.ш.ғ.д, проф., РГА корр-мүшесі, Ресей.

Г л а в н ы й р е д а к т о р

Есполов Т.И.,
доктор эконом. наук, проф.,
вице-президент и академик НАН РК

Р е д а к ц и о н на я кол л е г и я:

Байзаков С.Б., доктор эконом. наук, проф., академик НАН РК (заместитель главного редактора); **Тиреуов К.М.**, доктор эконом. наук., проф., академик НАН РК (заместитель главного редактора); **Елешев Р.Е.**, доктор техн. наук, проф., академик НАН РК; **Рай А.Г.**, доктор техн. наук, проф., академик НАН РК; **Иванов Н.П.**, доктор ветеринар. наук, проф., академик НАН РК; **Кешуов С.А.**, доктор техн. наук, проф., академик НАН РК; **Мелдебеков А.**, доктор сельхоз. наук, проф., академик НАН РК; **Чоманов У.Ч.**, доктор техн. наук, проф., академик НАН РК; **Елюбаев С.З.**, доктор сельхоз. наук, проф., академик НАН РК; **Садыкулов Т.**, доктор сельхоз. наук, проф., академик НАН РК; **Сансызыбай А.Р.**, доктор сельхоз. наук, проф., член-корр. НАН РК; **Умбетаев И.**, доктор сельхоз. наук, проф., академик НАН РК; **Оспанов С.Р.**, доктор сельхоз. наук, проф., Почетный член НАН РК; **Олейченко С.И.**, доктор сельхоз. наук, проф.; **Кененбаев С.Б.**, доктор сельхоз. наук, проф., член-корр. НАН РК; **Омбаев А.М.**, доктор сельхоз. наук, проф член-корр. НАН РК; **Молдашев А.Б.**, доктор эконом. наук, проф., Почетный член НАН РК; **Сагитов А.О.**, доктор биол. наук, академик НАН РК; **Сапаров А.С.**, доктор сельхоз. наук, проф., академик АСХН РК; **Балгабаев Н.Н.**, доктор сельхоз. наук, проф.; **Умирзаков С.И.**, доктор техн. наук, проф.; **Султанов А.А.**, доктор ветеринар. наук, проф., академик АСХН РК; **Алимкулов Ж.С.**, доктор техн. наук, проф., академик АСХН РК; **Сарсембаева Н.Б.**, доктор ветеринар. наук, проф.

Р е д а к ц и о н н ы й с о в е т:

Fasler-Kan Elizaveta, Dr., University of asel Switzeland; **Koolmees Petrus Adrianus**, Prof. Dr., Utrecht University, The Netherlands; **Babadoost-Kondri Mohammad**, Prof., University of Illinois, USA; **Yus Aniza Binti Yusof**, Dr., University Putra, Malayzia; **Hesseln Hayley Fawn**, As.Prof., University of Saskatchewan, Canada; **Alex Morgounov**, Pr., International Maize and Wheat Improvement Center Turkey; **Андреш С.**, академик НАН Республики Молдова; **Гаврилюк Н.Н.**, академик НАН Украины; **Герасимович Л.С.**, академик НАН Республики Беларусь; **Мамедов Г.**, академик НАН Республики Азербайджан; **Шейко И.П.**, академик НАН Республики Беларусь; **Жалнин Э.В.**, доктор техн. наук, проф., Россия; **Боинчан Б.**, доктор сельхоз. наук, проф., Республика Молдова; **Юлдашбаев Ю.А.**, доктор сельхоз. наук, проф., член-корр. РАН, Россия.

Известия Национальной академии наук Республики Казахстан. Серия аграрных наук.

ISSN 2224-526X

Собственник: РОО «Национальная академия наук Республики Казахстан» (г. Алматы)

Свидетельство о постановке на учет периодического печатного издания в Комитете информации и архивов Министерства культуры и информации Республики Казахстан № 10895-Ж, выданное 30.04.2010 г.

Периодичность 6 раз в год

Тираж: 300 экземпляров

Адрес редакции: 050010, г. Алматы, ул. Шевченко, 28, ком. 219-220, тел. 272-13-19, 272-13-18

<http://agricultural.kz/index.php/en/>

© Национальная академия наук Республики Казахстан, 2019

Адрес типографии: ИП «Аруна», г. Алматы, ул. Муратбаева, 75

Chief Editor

Espolov T.I.

Dr. economy. Sciences, prof.,
Vice President and academician of the NAS RK

Editorial Board:

Baizakov S.B., Dr. of economy sciences, prof., academician of NAS RK (deputy editor); **Tireuov K.M.**, Doctor of Economy Sciences., prof., academician of NAS RK (deputy editor); **Eleshev R.E.**, Dr. Of agricultural sciences, prof., academician of NAS RK; **Rau A.G.**, Dr. sciences, prof., academician of NAS RK; **Ivanov N.P.**, Dr. of veterinary sciences, prof., academician of NAS RK; **Keshuov S.A.**, Dr. sciences, prof., academician of NAS RK; **Meldebekov A.**, doctor of agricultural sciences, prof., academician of NAS RK; **Chomanov U.Ch.**, Dr. sciences, prof., academician of NAS RK; **Yelyubayev S.Z.**, Dr. of agricultural sciences, prof., academician of NAS RK; **Sadykulov T.**, Dr. Farm. Sciences, prof., academician of NAS RK; **Baimukanov D.A.**, doctor of agricultural sciences, prof., corresponding member NAS RK; **Sansyzbai A.R.**, doctor of agricultural sciences, prof., corresponding member NAS RK; **Umbetaev I.**, Dr. Farm. Sciences, prof., academician of NAS RK; **Ospanov S.R.**, Dr. agricultural sciences, prof., Honorary Member of NAS RK; **Olychenko S.N.**, Dr. Of agricultural sciences, prof.; **Kenenbayev S.B.**, Dr. Agricultural sciences, prof., corresponding member NAS RK; **Ombayev A.M.**, Dr. Agricultural sciences, Prof. corresponding member NAS RK; **Moldashev A.B.**, Doctor of Economy sciences, prof., Honorary Member of NAS RK; **Sagitov A.O.**, Dr. biol. sciences, academician of NAS RK; **Saparov A.S.**, Doctor of agricultural sciences, prof., academician of NAS RK; **Balgabaev N.N.**, the doctor agricultural sciences, Prof.; **Umirzakov S.I.**, Dr. Sci. Sciences, Prof.; **Sultanov A.A.**, Dr. of veterinary sciences, prof., academician of the Academy of Agricultural Sciences of Kazakhstan; **Alimkulov J.C.**, Dr. of tekhnical sciences, prof., academician of the Academy of Agricultural sciences of Kazakhstan; **Sarsembayeva N.B.**, Dr. veterinary sciences, prof.

Editorial Board:

Fasler-Kan Elizaveta, Dr., University of Basel Switzeland; **Koolmees Petrus Adrianus**, Prof. Dr., Utrecht University, The Netherlands; **Babadoost-Kondri Mohammad**, Prof., University of Illinois, USA; **Yus Aniza Binti Yusof**, Dr., University Putra, Malayzia; **Hesseln Hayley Fawn**, As. Prof., University of Saskatchewan, Canada; **Alex Morgounov**, candidate of agricultural sciences, International Maize and Wheat Improvement Center Turkey; **Andresh S.**, academician of NAS of Moldova; **Gavriluk N.N.**, academician of NAS of Ukraine; **Gerasimovich L.S.**, academician of NAS of Belorussia; **Mamadov G.**, academician of NAS of Azerbaijan; **Sheiko I.P.**, academician of NAS of Belorussia; **Zhalnin E.V.**, Dr. of technical sciences, professor, Russia, **Boinchan B.**, doctor of agricultural sciences, prof., Moldova; **Yuldasbayev Y.A.**, doctor of agricultural sciences, prof., corresponding member of RAS, Russia.

News of the National Academy of Sciences of the Republic of Kazakhstan. Series of Agrarian Sciences.

ISSN 2224-526X

Owner: RPA "National Academy of Sciences of the Republic of Kazakhstan" (Almaty)

The certificate of registration of a periodic printed publication in the Committee of Information and Archives of the Ministry of Culture and Information of the Republic of Kazakhstan N 10895-Ж, issued 30.04.2010

Periodicity: 6 times a year

Circulation: 300 copies

Editorial address: 28, Shevchenko str., of.219-220, Almaty, 050010, tel. 272-13-19, 272-13-18,
<http://nauka-nanrk.kz> / agricultural.kz

© National Academy of Sciences of the Republic of Kazakhstan, 2019

Address of printing house: ST "Aruna", 75, Muratbayev str, Almaty

NEWS

OF THE NATIONAL ACADEMY OF SCIENCES OF THE REPUBLIC OF KAZAKHSTAN

SERIES OF AGRICULTURAL SCIENCES

ISSN 2224-526X

Volume 5, Number 53 (2019), 45 – 53

<https://doi.org/10.32014/2019.2224-526X.59>

UDC 619: 595.2. 632.9

**I. I. Temreshev¹, A. M. Uspanov¹, A. B. Yeszhanov²,
A. M. Kenzhegaliev¹, A. M. Makezhanov¹, B. B. Bolatbekova¹**

¹LLP Zh. Zhiembaev Kazakh Research Institute of Plant Protection and Quarantine,
Ministry of Agriculture of Republic of Kazakhstan, Almaty, Kazakhstan,

²M. Aikimbaev Kazakh Scientific Center for Quarantine and Zoonotic Diseases, Almaty, Kazakhstan.
E-mail: temreshev76@mail.ru, u_alibek@mail.ru, aidyn.eszhanov@gmail.com, arnur_1992@mail.ru,
Makezhanov.arman@bk.ru, bolatbekova_beikhan@mail.ru

**ABOUT THE RESULTS OF LABORATORY TESTS
OF THE BIOLOGICAL DRUG ACTHAROPHYT
ON DIFFERENT SPECIES OF ARTHROPOD PESTS**

Abstract. The article presents data on laboratory testing of the efficacy of the Actarofit, a new biotechnological insectoacaricide of contact action, from a group of avermectins against several species of pest arthropods from different orders - an imagines of a painted bug *Eurydemaornata*(Linnaeus, 1758) (Hemiptera, Heteroptera, Pentatomidae), great green bush-cricketlarvae *Tettigoniaviridissima*(Linnaeus, 1758) (Orthoptera, Tettigoniidae), larvae of *Gomphomastaxclavata* (Ostromov, 1881) (Orthoptera, Eumastacidae)and *Chorthippusapricarius* (Linnaeus, 1758) (Orthoptera, Acrididae),imago of healer-beetle *Ulomoidesdermestoides*(Fairmaire, 1893) (Coleoptera, Tenebrionidae), wax moth caterpillars *Galleria mellonella*(Linnaeus, 1758) (Lepidoptera, Pyralidae), as well as blood sucking ixodid tick *Haemaphysalis punctata*Canestrini and Fanzago, 1877 (Acari, Ixodida, Ixodidae).The most resistant to the effects of Actarofit among the tested pest arthropods are the painted bug *Eurydemaornata* (Linnaeus, 1758), the wax moth caterpillars *Galleria mellonella* (Linnaeus, 1758) and the healer-beetle *Ulomoidesdermestoides* (Fairmaire, 1893). The larvae of the great green bush-cricket *Tettigoniaviridissima*(Linnaeus, 1758), *Gomphomastaxclavata* (Ostromov, 1881) and *Chorthippusapricarius* (Linnaeus, 1758), as well as the imago of the blood-sucking ixodid tick, *Haemaphysalis punctata*Canestrini and Fanzago, 1877, were the most susceptible to drug effect. Possibly this difference is largely linked with the thickness of the chitinous cover of the studied objects, which to some extent protects them from the effects of Actarofit. Based on the obtained results, it can be said that the biological drug Actarofit may be used for treatment against the wax moth, but with measures to prevent application of the drug on imago and larvae of bees. It can also be used against various pests of agriculture, and some bloodsucking arthropods. But it should always be taken into account that the sensitivity of the pest object to the drug action.

Key words: Aktarofit, avermectins, biological drug, pest arthropods, insects, blood-sucking ticks, protective measures, ecologization, agriculture.

Introduction. Modern methods of pest control in agriculture are still widely used chemical insecticides. This is mainly due to the fact that the widespread use of chemical insecticides due to economic necessity, and today biological control methods, are less effective comparing to chemical means of protection. High impact speed and efficacy are their advantage. Despite this, pest insects are often able to produce resistance to insecticides, which is expressed in changes in cover permeability, detoxification and changes in sensitivity to insecticides. The undoubted disadvantage of chemical insecticides is also the impact on non-target species of arthropods, as well as accumulation in the soil, plant and animal tissues. As a result of the increasing use of chemicals in plant protection, an enormous amount of various chemical compounds enter the environment. There is evidence that pesticides used in agriculture are able to act as mutagens, causing cytotoxic and negative genetic effects [1-3, 7-9, 22]. Biological drugs, in

particular, neurotoxic compounds such as avermectins, can serve as an alternative means of environmentally friendly plant protection. Avermectins - antibiotics, insecticides, acaricides and nematicides with a 16-membered macrocyclic lactone ring, referring to macrolides, are a product of the vital activity of actinomycetes *Streptomyces avermitilis* (ex Burget al., 1979) Kim and Goodfellow, 2002 (syn. *Streptomyces avermectinius* Takahashi et al., 2002).

In 1984, their synthetic forms were obtained. The principle of action is that they stimulate the release of gamma-aminobutyric acid (GABA) from the nerve endings and increase the connection of GABA with receptor sites on the postsynaptic membrane of muscle cells. This leads to blocking the transmission of nerve impulses, resulting in paralysis and death. The effectiveness of avermectins is very dependent on temperature, for example, when it decreases from 24 to 17°C, toxicity to spider mites decreases by 7.6 times, and when increased to 32°C, it increases by 4.8 times. A big advantage of avermectins is their low toxicity to mammals and humans, non-toxicity to earthworms, as well as to plants, rapid decomposition under the action of sunlight without the formation of toxic residues. Avermectins are not absorbed by the roots of plants from the soil and do not accumulate in them [10-21, 23, 24]. The authors asked about the possibility of using avermectins against various pests of agriculture in the Republic of Kazakhstan. For this purpose, Aktarofit was chosen - a new biotechnological insectoacaricide of contact action for the destruction of pests of agricultural, ornamental, forest and fruit crops (produced by "Enzim", Ukraine). The active basis of the drug is a complex of natural avermectins of groups B1 and B2, at a concentration of 0.2%. The introduction of such drugs is fully consistent with the tasks of transfer and technology adaptation and improving the environmental friendliness and competitiveness of Kazakhstan's agricultural products, set by the President of the Republic of Kazakhstan N.A. Nazarbayev in his annual message to the people in 2018.

Material and methods. As a model object, we used various species of pest arthropods from different orders - imago of a painted-bug *Eurydemaornata* (Linnaeus, 1758) (Hemiptera, Heteroptera, Pentatomidae), great green bush-cricketlarvae *Tettigoniaviridissima* (Linnaeus, 1758) (Orthoptera, Tettigoniidae), berry grasshopper larvae *Gomphomastaxclavata* (Ostromov, 1881) (Orthoptera, Eumastacidae) and *Chorthippusapricarius* (Linnaeus, 1758) (Orthoptera, Acrididae), healer-beetle imago *Ulomoidesdermestoides* (Fairmaire, 1893) (Coleoptera, Tenebrionidae), great wax moth caterpillars *Galleria mellonella* (Linnaeus, 1758) (Lepidoptera, Pyralidae), as well as the bloodsucking ixodid tick *Haemaphysalis-punctata* Canestrini and Fanzago, 1877 (Acari, Ixodida, Ixodidae). All insects, except for the last two species, were captured in nature with the help of a butterfly net and manual collection by employees of LLP Zh. Zhiembaev KazRIPPQ. The healer-beetle and the great wax moth were artificially bred under laboratory conditions in the insect of the M. Aykimbaev KSCQZD. Accordingly, experiments on species taken in nature were made in the laboratory of biotechnology of LLP Zh. Zhiembaev KazRIPPQ, while over artificially bred - at the insectary of the M. Aykimbaev KSCQZD, where the drug was transferred according to the oral agreement between the performers. The drug was diluted by us in a concentration of 1%, at a ratio of 400 ml of ordinary running water with the addition of 4 ml of avermectin. After this, the solution was applied to the bottom of Petri dishes or a plastic glass with a volume of 1 liter for the experimental group, while for the control group water was used. Exposure for both experimental and control groups was 1, 2 and 5 minutes. The exposure for the control group was the same time with the addition of water. It should be noted that for each group of tested insects, 3 replications were carried out.

In order to calculate the average time after which the full immobilization of pests occurs upon contact with the drug, time was counted on a stopwatch, immediately after the insect was treated.

The treated insects were placed in a clean Petri dish or plastic cup and checked on the following days after contact with the drug, in case of any surviving individuals could be found.

Research results. The pests used for the experiments are shown in figures 1-7. Below (tables 1-6) are the results from which it is clear that 1% solution of the drug Actarofit has 100% effectiveness against all species of pest arthropods used in the laboratory experiment.



Figure 1 –
The painted bug *Eurydema ornata* (Linnaeus, 1758)



Figure 2 –
The larva of the great green bush-cricket
Tettigoniaviridissima (Linnaeus, 1758)



Figure 3 –
The larva of *Gomphomastax clavata* (Ostroumov, 1881)



Figure 4 –
The larva of *Chorthippus apicarius* (Linnaeus, 1758)



Figure 5 –
Healer-beetle *Uloiodes dermestoides* (Fairmaire, 1893)



Figure 6 –
Great wax moth *Galleria mellonella* (Linnaeus, 1758)



Figure 7 –
Bloodsucking ixodic tick *Haemaphysalis punctata* Canestrini
and Fanzago, 1877

As can be seen from the data in Table 1, the painted bug *Eurydema ornata* (Linnaeus, 1758) – a pest of cruciferous crops, died completely 20 minutes after the treatment with Aktarofit.

Table 1 – The results of the test of Acarofit on the painted bug *Eurydema ornata* (Linnaeus, 1758)

Date	Experiment					Control				
	3 min	7 min	10 min	15 min	20 min	3 min	7 min	10 min	15 min	20 min
16.05.2019	8	4	3	2	0	10	10	10	10	10
17.05.2019	0	0	0	0	0	10	10	10	10	10

100% of a great green bush-cricket larvae *Tettigoniaviridissima* (Linnaeus, 1758) a multiple eating pest died 15 minutes after treatment.

Table 2 – The results of the test of Aktarofit on the great green bush-cricketlarvae *Tettigoniaviridissima* (Linnaeus, 1758)

Date	Experiment					Control				
	3 min	7 min	10 min	15 min	20 min	3 min	7 min	10 min	15 min	20 min
16.05.2019	6	4	2	1	0	10	10	10	10	10
17.05.2019	0	0	0	0	0	10	10	10	10	10

Larvae of berry grasshopper *Gomphomastaxclavata* (Ostroumov, 1881) and *Chorthippusapricarius* (Linnaeus, 1758) (Orthoptera, Acrididae) - a pest of fruit crops and medicinal plants, were even less resistant than the previous 2 insect species. Their 100% death occurred 10 minutes after the treatment with Aktarofit (table 3).

Table 3 – Results of the test of Aktarofit on the *Gomphomastaxclavata* larvae (Ostroumov, 1881) and *Chorthippusapricarius* (Linnaeus, 1758) (Orthoptera, Acrididae)

Date	Experiment			Control		
	3 min	7 min	10 min	3 min	7 min	10 min
16.05.2019	4	2	0	10	10	10
17.05.2019	0	0	0	10	10	10

Data on the efficacy of Acarofit against the wax moth - a pest of beekeeping [4], and the healer-beetle, a pest of stocks, are given in tables 4, 5. We found that the average time beforethe total immobilization of the wax moth larvae was 228.02 seconds, the maximum 405.82 seconds, and a minimum 104.26 seconds. While the same indicator for the imago of the healer-beetle was 36.86 seconds, 68.18 seconds, and 25.04 seconds respectively. Differences were found in the reactivity of both species for the drug, it was found that the imago beetles had the stage of excitation almost immediately after they were in contact with the drug, while the wax moth larvae reacted much later (about 1 minute). Probably, the differences in sensitivity and time before the total immobilization occurs consisted in the species specific resistance of insects, as well as their body weight. However, this aspect was not studied by us in this study.

Table 4 – Results of the test of Avermectin on the great wax moth caterpillars *Galleria mellonella* (Linnaeus, 1758)

Date	Experiment			Control		
	1 minute	2 minutes	5 minutes	1 minute	2 minutes	5 minutes
16.04.2019	3	3	3	3	3	3
17.04.2019	0	0	0	3	3	3
	Experiment			Control		
	1 minute	2 minutes	5 minutes	1 minute	2 minutes	5 minutes
17.04.2019	3	3	3	3	3	3
18.04.2019	0	0	0	3	3	3
	Experiment			Control		
	1 minute	2 minutes	5 minutes	1 minute	2 minutes	5 minutes
18.04.2019	3	3	3	3	3	3
19.04.2019	0	0	0	3	3	3

Table 5 – The results of the test of Aktarofit on the imago of the healer-beetle *Uloiodesdermestoides* (Fairmaire, 1893)

Date	Experiment			Control		
	1 minute	2 minutes	5 minutes	1 minute	2 minutes	5 minutes
16.04.2019	5	5	5	5	5	5
17.04.2019	0	0	0	5	5	5
	Experiment			Control		
	1 minute	2 minutes	5 minutes	1 minute	2 minutes	5 minutes
17.04.2019	5	5	5	5	5	5
18.04.2019	0	0	0	5	5	5
	Experiment			Control		
	1 minute	2 minutes	5 minutes	1 minute	2 minutes	5 minutes
18.04.2019	5	5	5	5	5	5
19.04.2019	0	0	0	5	5	5

100% of blood-sucking ixodic tick *Haemaphysalis punctata* Canestrini and Fanzago, 1877 died 10 minutes after treatment with Actarofit(Table 6).

Table 6 – The results of the test of Aktarofit
on the blood-sucking ixodic tick *Haemaphysalis punctata* Canestrini and Fanzago, 1877

Date	Experiment			Control		
	3 min	7 min	10 min	3 min	7 min	10 min
16.05.2019	2	1	0	3	3	3
17.05.2019	0	0	0	0	0	0

Discussion of research results. As can be seen from the obtained results, the painted bug *Euryde-maornata* (Linnaeus, 1758), the great wax mothcaterpillar *Galleria mellonella* (Linnaeus, 1758) and the healer beetle *Uloiodesdermestoides* (Fairmaire, 1893) have the greatest resistance to the effects of Actarofitfrom tested pest arthropods. The larvae of the great green bush-cricket *Tettigoniaviridissima* (Linnaeus, 1758), *Gomphomastaxclavata* (Ostromov, 1881) and *Chorthippusapricarius* (Linnaeus, 1758), as well as the imago of the blood-sucking ixodic tick *Haemaphysalis punctata* Canestrini and Fanzago, 1877 proved to be most susceptible to the effects of the drug. It can be assumed that this difference is largely due to the thickness of the chitinous cover of the studied objects, which to some extent protects them from the effects of Actarofit. Based on the results obtained, it can be said that the biological drug Actarofit may be used for treatment against the wax moth, but with measures to prevent the ingestion of the drug on the imago and the larvae of bees. Also, it can be used against various pests of agriculture, and some bloodsucking arthropods. But, the sensitivity of the harmful object to the action of the drugshould be taken into account.

Findings. Thus, the drug from the group of avermectins Actarofit can be used against various pests, for example, true bags, which are found quite a lot on food crops [5, 6]. In addition, the high efficacy of this biological drug against the blood-sucking ixodic tick, *Haemaphysalis punctata* Canestrini and Fanzago, 1877, is of great interest. This species is a grazing three host parasite. Imago feed on mammals (several dozen species), larvae and nymphs can also feed on birds and reptiles. The vector of causative agents of Qfever, Crimean hemorrhagic fever, tick-borne encephalitis, tularemia, Lyme disease, etc., which is of great sanitary and epidemiological importance. Therefore, the effect of Aktarofit on this tick is important not only in the field of plant protection, but also to ensure epidemiological well-being.

Source of research funding. The work was prepared as part of the implementation of the projects of the Ministry of Agriculture of the Republic of Kazakhstan BR 06249249 "Development of an integrated system to increase productivity and improvement of farm animals breeding qualities, using the example of BayserkeAgro LLP" under subproject 2 "Improving the technology of cultivation and harvesting of forage

"crops" and BR 06249206 "Transfer, adaptation and the introduction of advanced technologies for the control of quarantine and especially dangerous pests to ensure the phytosanitary safety of the agro-industrial complex of the Republic of Kazakhstan" for task 3 "Biological control of herd species of pest locusts using GIS technologies".

**И. И. Темрешев¹, А. М. Успанов¹, А. Б. Есжанов²,
А. М. Кенжегалиев¹, А. М. Макежанов¹, Б. Б. Болатбекова¹**

¹ҚР АШМ ТОО Ж. Жилембаев атындағы

Қазак өсімдік корғау және карантин ғылыми зерттеу институты, Алматы, Қазақстан,

²М. Айқымбаев атындағы Қазақ зооноздық және карантиндік инфекциялар ғылыми орталығы,
Алматы, Қазақстан

**АКТАРОФИТ БИОЛОГИЯЛЫҚ ПРЕПАРАТЫНЫң
ӘРТҮРЛІ ЗИЯНДЫ АРТРОПОДТАРҒА АРНАЛҒАН
ЗЕРТХАНАЛЫҚ ЗЕРТТЕУЛЕРИНІң НӘТИЖЕЛЕРІ ТУРАЛЫ**

Аннотация. Мақалада әр түрлі отрядтардан зиянды буынаяқтылардың бірнеше түріне қарсы авермектиндер тобынан контакттілі әсер ететін жаңа биотехнологиялық инсектоакарицид - Актарофит тиімділігін зертханалық тестілеу бойынша деректер келтіріледі – қандала *Eurydema ornata* (Linnaeus, 1758) (Hemiptera, Heteroptera, Pentatomidae), *Tettigonia viridissima* (Linnaeus, 1758) (Orthoptera, Tettigoniidae), *Gomphomastax clavata* (Ostroumov, 1881) (Orthoptera, Eumastacidae) және қоңыр саяқ шегіртке *Chorthippus apricarius* (Linnaeus, 1758) (Orthoptera, Acrididae), *Uloimoides dermestoides* (Fairmaire, 1893) (Coleoptera, tenebrionidae), *Galleria mellonella* (Linnaeus, 1758) (Lepidoptera, Pyralidae), сондай-ақ қан сорғыш иксод кенесінде *Haemaphysalis punctata* Canestrini and Fanzago, 1877 (Acari, Ixodida, Ixodidae). Сыналған зиянды буынаяқтардан Актарофит әсеріне ең жоғары тәзімділікке боялған қандала *Eurydema ornata* (Linnaeus, 1758), *Galleria mellonella* (Linnaeus, 1758) және *Uloimoides dermestoides* (fairmaire, 1893) балауыздары ие. Құрттар жасыл шекшек *Tettigonia viridissima* (Linnaeus, 1758), жидек биелер *Gomphomastax clavata* (Ostroumov, 1881) және қоңыр керек *Chorthippus apricarius* (Linnaeus, 1758), сондай-ақ имаго кровососущего иксодового кене *Haemaphysalis punctata* Canestrini and Fanzago, 1877 қалсаңыз, ең сезімтал әсеріне препарат. Мұндай айырмашылық көбінесе зерттелетін объектілердің хитин жабынының қалындығына байланысты болуы мүмкін, ол оларды Актарофиттің әсерінен корғайды. Алынған нәтижелерді негізге ала отырып, биологиялық Актарофит препаратын балауыз балына қарсы өңдеу үшін пайдалануға болады, бірақ препараттың имаго мен бал араларының личинкаларына тұсуінің алдын алу жөніндегі шаралармен бірге. Сондай-ақ оны ауыл шаруашылығының әртүрлі зиянкестеріне және кейбір қан соратын буынаяқтыларға қарсы қолдануға болады. Бірақ бұл ретте зиянды объектінің препараттың әсеріне сезімталдығын міндетті түрде ескеру керек.

Түйін сөздер: Актарофит, авермектиндер, биологиялық препарат, зиянды буынаяқтылар, жәндіктер, қан соратын кенелер, корғау шаралары, экологизация, ауыл шаруашылығы.

**И. И. Темрешев¹, А. М. Успанов¹, А. Б. Есжанов²,
А. М. Кенжегалиев¹, А. М. Макежанов¹, Б. Б. Болатбекова¹**

¹ТОО Казахский научно-исследовательский институт защиты и карантина растений
им. Ж. Жилембаева МСХ РК, Алматы, Казахстан,

²Казахский научный центр зоонозных и карантинных инфекций им. М. Айкимбаева, Алматы, Казахстан

**О РЕЗУЛЬТАТАХ ЛАБОРАТОРНЫХ ИСПЫТАНИЙ БИОЛОГИЧЕСКОГО ПРЕПАРАТА
АКТАРОФИТ НА РАЗЛИЧНЫХ ВИДАХ ВРЕДНЫХ ЧЛЕНИСТОНОГИХ**

Аннотация. В статье приводятся данные по лабораторному тестированию эффективности Актарофита – нового биотехнологического инсектоакарицида контактного действия из группы авермектинов против нескольких видов вредных членистоногих из разных отрядов – имаго разукрашенного клопа *Eurydema ornata* (Linnaeus, 1758) (Hemiptera, Heteroptera, Pentatomidae), личинок зеленого кузнецика *Tettigonia viridissima* (Linnaeus, 1758) (Orthoptera, Tettigoniidae), личинок ягодной кобылки *Gomphomastax clavata* (Ostroumov, 1881) (Orthoptera, Eumastacidae) и бурого конька *Chorthippus apricarius* (Linnaeus, 1758) (Orthoptera, Acrididae), имаго чернотелки жука-знахаря *Uloimoides dermestoides* (Fairmaire, 1893) (Coleoptera, Tenebrionidae),

гусениц большой восковой моли *Galleriamellonella* (Linnaeus, 1758) (Lepidoptera, Pyralidae), а также на кровососущем иксодовом клеще *Haemaphysalis punctata* Canestriniand Fanzago, 1877 (Acari, Ixodida, Ixodidae) наибольшей устойчивостью к воздействию Актарофита из протестируемых вредных членистоногих обладают разукрашенный клоп *Eurydema ornata* (Linnaeus, 1758), гусеницы большой восковой моли *Galleriamellonella* (Linnaeus, 1758) и жук-знахарь *Uloiodes dermestoides* (Fairmaire, 1893). Личинки зеленого кузнецика *Tettigoniaviridissima* (Linnaeus, 1758), ягодной кобылки *Gomphomastax clavata* (Ostromov, 1881) и бурого конька *Chorthippus apricarius* (Linnaeus, 1758), а также имаго кровососущего иксодового клеща *Haemaphysalis punctata* Canestriniand Fanzago, 1877 оказались наиболее восприимчивы к воздействию препарата. Возможно, что подобное различие во многом связано с толщиной хитинового покрова исследуемых объектов, которое в какой-то мере защищает их от воздействия Актарофита. Исходя из полученных результатов, можно говорить о том, что биологический препарат Актарофит возможно использовать для обработки против восковой моли, однако с мерами по предупреждению попадания препарата на имаго и личинок пчел. Также его можно применять против различных вредителей сельского хозяйства, и некоторых кровососущих членистоногих. Но при этом следует обязательно учитывать чувствительность вредного объекта к действию препарата.

Ключевые слова: Актарофит, авермектины, биологический препарат, вредные членистоногие, насекомые, кровососущие клещи, защитные мероприятия, экологизация, сельское хозяйство.

Information about authors:

Temreshev I. I., LLP Zh. Zhiembaev Kazakh Research Institute of Plant Protection and Quarantine, Ministry of Agriculture of Republic of Kazakhstan, Almaty, Kazakhstan; temreshev76@mail.ru; <https://orcid.org/0000-0003-0004-4399>

Uspanov A. M., LLP Zh. Zhiembaev Kazakh Research Institute of Plant Protection and Quarantine, Ministry of Agriculture of Republic of Kazakhstan, Almaty, Kazakhstan; u_alibek@mail.ru; <https://orcid.org/0000-0001-7122-8596>

Yeszhanov A. B., M. Aikimbaev Kazakh Scientific Center for Quarantine and Zoonotic Diseases, Almaty, Kazakhstan; aidyn.eszhanov@gmail.com; <https://orcid.org/0000-0001-6572-5668>

Kenzhegaliev A. M., LLP Zh. Zhiembaev Kazakh Research Institute of Plant Protection and Quarantine, Ministry of Agriculture of Republic of Kazakhstan, Almaty, Kazakhstan; arnur_1992@mail.ru; <https://orcid.org/0000-0002-0308-222X>

Makezhanov A. M., LLP Zh. Zhiembaev Kazakh Research Institute of Plant Protection and Quarantine, Ministry of Agriculture of Republic of Kazakhstan, Almaty, Kazakhstan; Makezhanov.arman@bk.ru; <https://orcid.org/0000-0003-1587-1979>

Bolatbekova B. B., LLP Zh. Zhiembaev Kazakh Research Institute of Plant Protection and Quarantine, Ministry of Agriculture of Republic of Kazakhstan, Almaty, Kazakhstan; bolatbekova_beikhan@mail.ru; <https://orcid.org/0000-0001-5924-493X>

REFERENCES

- [1] Barilyak N.R., Buzhievskaia T.I., Bykorez A.I. (1989). Genetic effects of environmental pollution. Kiev: Naukova Dumka (in Russ.).
- [2] Begimbetova D.A., Kolumbaeva S.Zh., Kalimagambetov A.M. (2009). The genotoxic effect of fipronil on rats of different ages // Experimental Biology. 40. 1: 96-99 (in Russ.).
- [3] Benkovskaya G.V., Leontyeva T.L., Udalov M.B. (2008). Resistance of the Colorado potato beetle to insecticides in the Southern Urals // Agrochemistry. 8: 55-59 (in Russ.).
- [4] Grobov O.F., Smirnov A.M., Popov E.T. (1987). Diseases and pests of honey bees. M.: Agropromizdat (in Russ.).
- [5] Esenbekova P.A., Temreshev I.I., Kenzhegaliev A.M., Tursynkulov A.M., Dosmukhambetov T.M. (2019). Hemiptera (Hemiptera: Heteroptera) – pest and grain (barley, triticale, wheat) Bayserke-Agro LLP // News of the National academy of sciences and the Republic of Kazakhstan. Series of agricultural sciences. 2 (50): 21-30. <https://doi.org/10.32014/2019.2224-526X.12> ISSN 2224-526X
- [6] Esenbekova P.A., Temreshev I.I., Kenzhegaliev A.M., Tursynkulov A.M., Dosmukhambetov T.M. (2019). Bugs (Hemiptera: Heteroptera) – pests of alfalfa "Bayserke-Agro" LLP // News of the National academy of sciences of the Republic of Kazakhstan. Series of agricultural sciences. 2 (50): 55-65. <https://doi.org/10.32014/2019.2224-526X.17> ISSN 2224-526X (in Russ.).
- [7] Korotkova L.I. (2007). Accumulation of organochlorine compounds in the commercial fish of the north-eastern part of the Black Sea. Ichthyological studies in inland waters. Materials of the International Scientific Conference. Saransk: Mordovia State University. P. 91-93 (in Russ.).

- [8] Slivinsky G.G., Temreshev I.I., Isenova G.D., Kozhabayeva G.E. (2016). Aquatic beetles as bioindicators of environmental conditions in the water bodies of South Kazakhstan // *ActaBiologica Sibirica.* 2 (3): 46-53.<http://dx.doi.org/10.14258/abs.v2i3.1454> (in Russ.).
- [9] Smirnov A.A. (2006). Screening of insectoacaricides and their mixtures, creating a compound with a synergistic effect // *Veterinary Pathology.* 1: 118-121 (in Russ.).
- [10] Roslavitseva S.A. (1987). New group of insectoacaricides and nematocides // *Agrochemistry.* 7: 130 (in Russ.).
- [11] Roslavitseva S.A., Didenko L.N. (2007). New in the problem of insect acaricide resistance of arthropods // *Agrochemistry.* 7: 88-91 (in Russ.).
- [12] Burg R.W., Miller B.M., Baker E.E., Birnbaum J., Currie S.A., Hartman R., Kong Y.-L., Monaghan R.L., Olson G., Putter I., Tunac J.B., Wallick H., Stapley E.O., Oiwa R., Omura S. (1979). Avermectins, New Family of Potent Anthelmintic Agents: Producing Organism and Fermentation // *Antimicrobial Agents and Chemotherapy.* 15(3): 361-7. doi:10.1128/AAC.15.3.361.
- [13] Clark J.K., Scott J.G., Campos F., Bloomquist J.R. (1995). Resistance to Avermectins: Extent, Mechanisms, and Management Implications // *Annual Review of Entomology.* 40: 1-30. doi:10.1146/annurev.en.40.010195.000245.
- [14] Cully Doris F., Vassilatis Demetrios K., Liu Ken K., Paress Philip S., Van Der Ploeg, Lex H. T., Schaeffer James M., Arena Joseph P. (1994). Cloning of an avermectin-sensitive glutamate-gated chloride channel from *Caenorhabditiselegans* // *Nature.* 371 (6499). 707-11. Bibcode:1994Natur.371..707C. doi:10.1038/371707a0.
- [15] Gao Q., Tan G.-Y., Xia X., Zhang L. (2017). Learn from microbial intelligence for avermectin overproduction. *Current Opinion in Biotechnology.* 48.: 251-257. doi: 10.1016/j.copbio.2017.08.016.
- [16] Ikeda H., Nonomiya T., Usami M., Ohta T., Omura S. (1999). Organization of the biosynthetic gene cluster for the polyketide anthelmintic macrolide avermectin in *Streptomyces avermitilis* // *Proceedings of the National Academy of Sciences.* 96(17). P. 9509-9514. doi:10.1073/pnas.96.17.9509.
- [17] Jinsong Chen, Jin Miao, Mei Liu, Xueling Liu. (2016). Different fates of avermectin and artemisinin in China // *Science China. Life sciences.* 59 (6).doi: 10.1007/s11427-016-5065-y.
- [18] Ōmura Satoshi, ShiomiKazuro. (2007). Discovery, chemistry, and chemical biology of microbial products // *Pure and Applied Chemistry.* 79(4). 581-591. doi:10.1351/pac200779040581.
- [19] Pitterma T., Cassayre J., Hüter O.F., Jung Pierre M.J., Maienfisch P., Kessabi F.M., Quaranta L., Tobler H. (2009). New ventures in the chemistry of avermectins // *Bioorganic & Medicinal Chemistry.* 17 (12). 4085-4095. doi:10.1016/j.bmc.2008.12.069.
- [20] Siddique S., Syed Q., Adnan A., Nadeem M., Irfan M., Ashraf Qureshi F. (2013). Production of Avermectin B1b from *Streptomyces avermitilis* 41445 by Batch Submerged Fermentation // *Jundishapur J. Microbiol.* 6 (8): e7198. doi: 10.5812/jjm.7198.
- [21] Takahashi Y. (2002). *Streptomyces avermectinii sp. nov.*, an avermectin-producing strain // *International Journal of Systematic and Evolutionary Microbiology.* 52 (6): 2163-8. doi:10.1099/ijss.0.02237-0.
- [22] Temreshev I.I., Esenbekova P.A., Sagitov A.O., Mukhamadiev N.S., Sarsenbaeva G.B., Ageenko A.V., Homziak J. (2018). Evaluation of the effect of locally produced biological pesticide (AkKobelek™) on biodiversity and abundance of beneficial insects in four forage crops in the Almaty region of Kazakhstan // *International Journal of Environment, Agriculture and Biotechnology (IJEAB).* 3. 1.: 72-91. <http://dx.doi.org/10.22161/ijeab/3.1.10>. ISSN: 2456-1878. Thomson Reuters Researcher ID: E-2759-2017.
- [23] Yoon Y.J., Kim E.-S., Hwang Y.-S., Choi C.-Y. (2004). Avermectin: Biochemical and molecular basis of its biosynthesis and regulation // *Applied Microbiology and Biotechnology.* 63 (6): 26-34. doi:10.1007/s00253-003-1491-4.
- [24] Zhang Changsheng, Albermann Christoph, Fu Xun, Thorson Jon S. (2006). The in Vitro Characterization of the Iterative Avermectin Glycosyltransferase AveBI Reveals Reaction Reversibility and Sugar Nucleotide Flexibility // *Journal of the American Chemical Society.* 128 (51).: 16420-1. doi:10.1021/ja065950.

Publication Ethics and Publication Malpractice in the journals of the National Academy of Sciences of the Republic of Kazakhstan

For information on Ethics in publishing and Ethical guidelines for journal publication see <http://www.elsevier.com/publishingethics> and <http://www.elsevier.com/journal-authors/ethics>.

Submission of an article to the National Academy of Sciences of the Republic of Kazakhstan implies that the described work has not been published previously (except in the form of an abstract or as part of a published lecture or academic thesis or as an electronic preprint, see <http://www.elsevier.com/postingpolicy>), that it is not under consideration for publication elsewhere, that its publication is approved by all authors and tacitly or explicitly by the responsible authorities where the work was carried out, and that, if accepted, it will not be published elsewhere in the same form, in English or in any other language, including electronically without the written consent of the copyright-holder. In particular, translations into English of papers already published in another language are not accepted.

No other forms of scientific misconduct are allowed, such as plagiarism, falsification, fraudulent data, incorrect interpretation of other works, incorrect citations, etc. The National Academy of Sciences of the Republic of Kazakhstan follows the Code of Conduct of the Committee on Publication Ethics (COPE), and follows the COPE Flowcharts for Resolving Cases of Suspected Misconduct (http://publicationethics.org/files/u2/New_Code.pdf). To verify originality, your article may be checked by the Cross Check originality detection service <http://www.elsevier.com/editors/plagdetect>.

The authors are obliged to participate in peer review process and be ready to provide corrections, clarifications, retractions and apologies when needed. All authors of a paper should have significantly contributed to the research.

The reviewers should provide objective judgments and should point out relevant published works which are not yet cited. Reviewed articles should be treated confidentially. The reviewers will be chosen in such a way that there is no conflict of interests with respect to the research, the authors and/or the research funders.

The editors have complete responsibility and authority to reject or accept a paper, and they will only accept a paper when reasonably certain. They will preserve anonymity of reviewers and promote publication of corrections, clarifications, retractions and apologies when needed. The acceptance of a paper automatically implies the copyright transfer to the National Academy of Sciences of the Republic of Kazakhstan.

The Editorial Board of the National Academy of Sciences of the Republic of Kazakhstan will monitor and safeguard publishing ethics.

Правила оформления статьи для публикации в журнале смотреть на сайте:

[www:nauka-nanrk.kz](http://nauka-nanrk.kz)

<http://agricultural.kz/index.php/en/>

Редактор М. С. Ахметова, Т. М. Апендиев, Д. С. Аленов
Верстка на компьютере Д. Н. Калкабековой

Подписано в печать 14.10.2019.
Формат 60x881/8. Бумага офсетная. Печать – ризограф.
8,2 п.л. Тираж 300. Заказ 5.